

- a ‘II’ indecates HII region; ‘H’ indicates optical jet or HH object; ‘2’ indicates H₂ jet; ‘W’ indicates water maser; ‘I’ indicates infall or collapse.
- b The maps of these sources are not complete or confused with other sources.
- c These sources are detected with CO J=3-2 or higher.
- d These outflows without corresponding young stellar objects.
- e The underlined R is quoted from references.
- f This source includes NGC7538 IRS1, IRS9, IRS11, and A.
- g The distance is one of the assumed values(cf. Ref. 123).
- h The mass is calculated under optical thick.
- i This reference proposes that 05358+3543 consists of CO flow A, B and a SiO flow C.
- j The authors of Ref. 326 proposed that this source has 2 overlapping outflows.
- k The HH objects detected before 1999 were all from Ref. 251.
- l This source includes RNO43S and RNO43N.
- m The parameters are derived from C¹⁸O observation.
- n This source includes IC1396N BIMA1 and BIMA2.
- o $\dot{M}_f$ is not used if it is deduced by M/τ .
- p It was not identified as bipolar outflow by Ref. 191.
- q Ref. 327 reported that there may be two outflows.
- r Its reshift lobe is probably masked by the outflow of L379 IRAS(3).
- s This timescale is calculated with the assumption that the clump has a tangential velocity of 200 km/s.
- t This source includes NGC2024 Fir 4, 5 and 6.
- u This source includes NGC7129 FIRS2-MM1, FIRS2-IR, FIRS1-IRS6 and FIRS1-MM1.
- v Parameters calculated here are not for the whole outflow but only for the n4 region.
- w The author of Ref. 354 proposed that outflows at this position are not driven by FIR sources but by jets from faint undetected yet low-mass stars; the parameters in Table 1a are the total of three outflows identified.

References

- 1, Bally & Lada, 1983;
- 2, Cantó et al., 1984;
- 3, Snell et al., 1990;
- 4, Brackmann & Scoville, 1980;
- 5, Arquilla & Goldsmith, 1984;
- 6, Nakano & Yoshida, 1986;
- 7, Lada, 1985;
- 8, Levreault, 1985;
- 9, Bachiller et al., 1990b;
- 10, Lada & Harvey, 1981;
- 11, Snell et al., 1984;
- 12, Levreault, 1988;
- 13, Mao et al., 1989;
- 14, Goldsmith et al., 1984;
- 15, Yamashita et al., 1989a;
- 16, Phillips et al., 1988;
- 17, Edwards & Snell, 1984;
- 18, Edwards & Snell, 1983;
- 19, Snell & Edwards, 1981;
- 20, Beichman et al., 1984;
- 21, Goldsmith et al., 1986;
- 22, Myers et al., 1988;
- 23, Myers et al., 1987;
- 24, Heyer et al., 1987;
- 25, Snell et al., 1980;
- 26, Strom et al., 1976;
- 27, Lichten, 1982a;
- 28, Beichman et al., 1986;
- 29, Liljeström et al., 1989;
- 30, Frerking & Langer, 1982;
- 31, Wu et al., 1992;
- 32, Snell et al., 1988;
- 33, Fukui et al., 1986;
- 34, Morgan & Bally 1991;
- 35, Ziurys et al., 1990;
- 36, Wouterloot & Brand, 1989;
- 37, Schmid-Burgk et al. 1989;
- 38, Erickson et al., 1982;
- 39, Kwan & Scoville 1976;
- 40, Olofsson et al., 1982;
- 41, Zuckerman et al., 1976;
- 42, Hilton et al., 1986;
- 43, Snell & Edwards, 1982;
- 44, Campbell & Thompson, 1984;
- 45, Bally, 1982;
- 46, Lichten, 1982b;
- 47, Margulis et al., 1988;
- 48, Haschick et al., 1981;
- 49, Rodriguez et al., 1982;
- 50, Blitz & Thaddeus, 1980;
- 51, Lada & Gautier, 1982;
- 52, Levreault, 1983;
- 53, Margulis & Lada 1986;
- 54, Dent et al., 1984;
- 55, Rodriguez et al., 1980;
- 56, Tamura et al., 1990;
- 57, Fischer et al., 1982;
- 58, Simon et al., 1984;
- 59, Mozurkewich et al., 1986;
- 60, Ho et al., 1982;
- 61, Armstrong & Winnewisser, 1989;
- 62, Edwards & Snell, 1982;
- 63, Little et al., 1988;
- 64, Lada et al., 1984;
- 65, Dickel et al., 1978;
- 66, Levreault 1984;
- 67, White & Phillips 1981;
- 68, Sato & Fukui 1989;
- 69, Mathieu et al., 1988;
- 70, Rodríguez et al., 1981;
- 71, Haikala & Laureijs, 1989;
- 72, Dent et al., 1985a;
- 73, Dent et al., 1985b;
- 74, Cantó et al., 1981;
- 75, Reipurth & Olberg, 1991;
- 76, Kameya et al., 1989;
- 77, Evans et al., 1981a;
- 78, Harvey et al., 1984;
- 79, Schwartz et al., 1983;
- 80, Calvet et al., 1983;
- 81, Wolf et al., 1990;
- 82, Hayashi et al., 1987;
- 83, Evans et al., 1981b;
- 84, Terebey et al., 1989;
- 85, Yang et al., 1990;
- 86, Fukui, 1989;
- 87, Arquilla & Kwok 1987;

- 88, Fischer et al., 1985;
- 89, Sugitani et al., 1989;
- 90, Casoli et al., 1986;
- 91, Schwartz et al., 1988;
- 92, Loren, 1989;
- 93, Parker et al., 1988;
- 94, Harvey & Forveille 1988;
- 95, Scoville et al., 1986;
- 96, Bachiller & Cernicharo, 1990;
- 97, Xie & Goldsmith, 1990;
- 98, Clark, 1986;
- 99, Wouterloot et al., 1989;
- 100, Vallée & Avery, 1990;
- 101, Sanders & Willner, 1985;
- 102, Richer, 1990;
- 103, Schmid-Burgk et al., 1990;
- 104, Yang et al., 1991;
- 105, Mizuno et al., 1990;
- 106, Andre et al., 1990;
- 107, Richer et al., 1989;
- 108, Schwartz, et al., 1985;
- 109, Davidson & Jaffe, 1984;
- 110, Iwata et al., 1988;
- 111, Olberg et al., 1992;
- 112, Mattila et al., 1989;
- 113, Smith & Fischer, 1992;
- 114, Reipurth, 1989;
- 115, Reipurth & Graham, 1991;
- 116, Wilking et al., 1990;
- 117, Heyer et al., 1989;
- 118, Evans et al., 1994;
- 119, Koo et al., 1994;
- 120, Phillips & Mampaso 1991;
- 121, Bachiller et al., 1991;
- 122, Umemoto et al., 1992;
- 123, Odenwald & Schwartz 1993;
- 124, Ladd et al., 1991;
- 125, Barsony, 1989;
- 126, Bachiller et al., 1990a;
- 127, Chernin & Masson, 1991;
- 128, Moriarty-Schieven & Wanner, 1991;
- 129, Wilson & Mauersberger 1991;
- 130, Knee, 1992;
- 131, Wootten & Loren, 1987;
- 132, Margulis et al., 1990;
- 133, Fukui et al., 1988;
- 134, Mendoza et al., 1990;
- 135, Liseau et al., 1988;
- 136, Walker et al., 1988;
- 137, Kutner et al., 1982;
- 138, Fukui et al., 1989;
- 139, Weintraub et al., 1993;
- 140, Loren et al., 1990;
- 141, Eiroa et al., 1992;
- 142, Gregório Hetem et al., 1990;
- 143, Cohen & Schwartz 1987;
- 144, Terebey et al., 1990;
- 145, Moriarty-Schieven et al., 1992;
- 146, Tamura et al., 1991;
- 147, Margulis et al., 1989;
- 148, Garden et al., 1991a;
- 149, Umemoto et al., 1991;
- 150, Duvert et al., 1990;
- 151, Koo, 1989;
- 152, Choi et al., 1993;
- 153, Margulis & Snell, 1989;
- 154, Avery et al., 1990;
- 155, Olberg et al., 1989;
- 156, Reipurth et al., 1993;
- 157, Armstrong, 1989;
- 158, Cohen 1980;
- 159, Kenyon et al., 1993;
- 160, Torrelles et al., 1993;
- 161, Morgan et al., 1991;
- 162, Dobashi et al., 1993;
- 163, Minchin et al., 1993;
- 164, Kastner et al., 1994;
- 165, Bally et al., 1994;
- 166, Hasegawa et al., 1994;
- 167, White, 1993;
- 168, Parker et al., 1991;
- 169, Blake et al., 1995;
- 170, Minchin et al., 1995;
- 171, Sato et al., 1994;
- 172, Jiang et al., 2001;
- 173, Zeilik et al., 1978;
- 174, Masson et al., 1990;
- 175, Chernin & Masson, 1992;

- 176, Haikala & Dietrich, 1989;
177, Bourke, 1995;
178, Yun & Clemens, 1992;
179, Wang et al., 1995;
180, Dobashi et al., 1994;
181, Wu et al., 1999a;
182, Xiang & Turner, 1995;
183, Wouterloot et al., 1988;
184, Moriarty-Schieven et al., 1995;
185, Yonekura et al., 1998;
186, Dobashi et al., 1998a;
187, Dobashi et al., 1998b;
188, Bourke et al., 1997;
189, Reipurth et al., 1997;
190, Bontemps et al., 1996a;
191, Wu et al., 2004;
192, Lyder et al., 1998;
193, Shepherd & Churchwell, 1996a;
194, Shepherd & Churchwell, 1996b;
195, Correia et al., 1997;
196, Wouterloot & Brand, 1999;
197, Chandler et al., 1996;
198, Wu et al., 1999b;
199, Yang & Wu, 1998;
200, Hirano et al., 1997;
201, Codella & Bachiller, 1999;
202, Hunter et al., 1997;
203, Mao, 1999;
204, Ladd & Hodapp, 1997;
205, Meyers-Rice & Lada, 1991;
206, Fich & Lada, 1997;
207, Zhang et al., 2001;
208, Henning et al., 2000a;
209, Gueth & Guilloteau, 1999;
210, Chen et al., 1997;
211, Langer et al., 1996;
212, Schreyer et al., 1997;
213, Shepherd et al., 2000;
214, Gueth et al., 1996;
215, Lee et al., 2000;
216, Benson et al., 1984;
217, Shepherd & Kurtz, 1998;
218, Sandell et al., 1999;
219, Henning et al., 2000b;
220, Yun et al., 2001;
221, Sugitani & Ogura, 1994;
222, Knee & Sandell, 2000;
223, Cabrit et al., 1988;
224, Langer et al., 1996;
225, Shepherd et al., 1997;
226, Magnier et al., 1999;
227, Reipurth, 1999b;
228, Cabrit & Bertout, 1992;
229, Bally et al., 1999;
230, Zhou et al., 1993;
231, Zhou, 1999;
232, Gregersen et al., 1997;
233, Walker et al., 1986;
234, Zhou, 1995;
235, Myers, et al., 1996;
236, Ward-Thompson et al., 1996;
237, Beuther et al., 2002a;
238, Arce & Goodman, 2001;
239, Ridge & Moore, 2001;
240, Yun & Clemens, 1994;
241, Hirano et al., 1998;
242, Wolf-Chase et al., 2000;
243, Monin et al., 1996;
244, Palacios & Eiroa, 1999;
245, Moriarty-Schieven & Snell, 1988;
246, Hogerheijde et al., 1998;
247, Hunter et al., 1994;
248, Bachiller et al., 1995a;
249, Chernin & Wright, 1996;
250, Kawamura et al., 1999;
251, Reipurth, 1999;
252, Yamashita et al., 1989b;
253, Myers et al., 1995;
254, Sugitani et al., 1991;
255, Churchwell, 1999;
256, Yamaguchi et al., 1999a;
257, Onishi et al., 1999;
258, Dobashi et al., 2001;
259, Yamaguchi et al., 1999b;
260, Saito et al., 1999;
261, Chernin & Masson, 1995;
262, Aso et al., 2000;
263, Moro-Martín et al., 1999;

- 264, Davis et al., 2000;
265, Davis et al., 1994a;
266, Eislöffel, 2000;
267, Hodapp & Ladd, 1995;
268, Davis et al., 1997;
269, Aspin, 1992;
270, Bachiller et al., 1994;
271, Hartigan et al., 1989;
272, Yu et al., 1999;
273, Stanke et al., 1998a;
274, Gomez et al. 1997;
275, Davis et al., 1994b;
276, Davis et al., 1995;
277, Yu et al., 1997;
278, Stanke et al., 2002;
279, Porras et al., 2000;
280, Davis et al., 2002;
281, Field et al., 1998;
282, Chen et al., 1999;
283, Luhman et al., 1997;
284, Chakraborty et al., 2000;
285, Massi et al., 1997;
286, Aspin, 1998;
287, Ryder et al., 1998;
288, Yun et al., 1997;
289, Heathcote et al., 1996;
290, Eislöffel et al., 2000;
291, Davis & Eislöffel, 1995;
292, Casali et al., 1993;
293, Davis et al., 1999a;
294, Aspin et al., 1991;
295, Holbrook, 1996;
296, Stanke et al., 2000;
297, Garden et al., 1990;
298, Poetzel et al., 1992;
299, Ayala et al., 1998;
300, Nisini et al., 2001;
301, Yu et al., 1995;
302, Davis et al., 1998;
303, Stanke et al., 1998b;
304, Cabrit et al., 1997;
305, Hartigan et al., 1995;
306, Kumar et al., 2002;
307, Shepherd et al., 1998;
308, Bloomer et al., 1998;
309, Longmore et al., 1986;
310, Velusamy & Langer, 1998;
311, Davis et al., 1996;
312, Mitchell & Hasegawa, 1991;
313, Hunter et al., 1995;
314, Beuther et al., 2002b
315, Qin & Wu, 2003a;
316, Yu et al., 2000;
317, Fuente et al., 2001;
318, Bourke, 2001;
319, Arce & Goodman, 2002;
320, Moreira et al., 2000;
321, Dobashi & Uehara, 2001;
322, White et al., 1997;
323, Sandell & Aspin, 1998;
324, Davis et al., 1999b;
325, Beuther et al., 2002c;
326, Aspin & Reipurth, 2000;
327, Kelly & Macdonald, 1996;
328, Ogura, Sugitani & Pickles, 2002;
329, Zhou et al., 1996;
330, Ohashi et al., 1997;
331, Choi et al., 1999;
332, Bally et al., 2002;
333, Yang et al., 1997;
334, Gregersen et al., 2000;
335, Lada & Fich, 1996;
336, Fich & Lada, 1998;
337, Kurtz et al., 1994;
338, Walsh et al., 1998;
339, McCutcheon et al., 1991;
340, Altenhoff et al., 1978;
341, Gregory & Condon, 1991;
342, Griffith et al., 1994;
343, Wright et al., 1994;
344, Wolf-Chase & Gregersen, 1997;
345, Comoretto et al., 1990;
346, Brand et al., 1994;
347, Valdetaro et al., 2001;
348, Mitchell et al., 1991;
349, Braz & Epchtein, 1983;
350, Scalise et al., 1989;
351, Bence et al., 1996;

- 352, Rodríguez & Canto, 1982
353, Moran & Rodríguez, 1980;
354, Mitchell, et al., 1995;
355, Hofner et al., 1996;
356, Mardones et al., 1997;
357, Olmi, Felli & Cesaroni, 1997;
358, Bence et al., 1998;
359, Molinari et al, 1998;
360, Chandler & Carlstrom, 1996;
361, Snell & Bally, 1986;
362, Garden et al., 1991b;
363, Cesaroni et al., 1997;
364, Richer et al., 1992;
365, White et al., 1995;
366, Tamura et al, 1996;
367, André et al., 1999;
368, Belloche et al., 2002;
369, Wootten et al., 1989.
370, Gueth et al., 2001;
371, Magnier et al., 1995;
372, Mao et al., 2002;
373, Beltrán et al., 2002;
374, Anglada & Rodríguez, 2002;
375, Girart et al., 2002;
376, Zhao et al., 1999;
377, Hilton & Lahulla, 1993;
378, Kamazaki et al., 2003;
379, Gibb & Little, 2000;
380, Shi et al., 2003;
381, Yu et al., 2003a;
382, Zhao et al., 2003;
383, Lee et al., 2002;
384, Phillips & Mampaso, 1996;
385, Dent & Aspin, 1992;
386, Reipurth et al, 2000;
387, Bontemps et al., 1996b;
388, Chernin, 1996;
389, Lefloch, 1996;
390, Bally et al., 1996a;
391, Bally et al., 1996b;
392, Noriega et al., 2002;
393, Patel et al., 2000;
394, Khanzadyan et al., 2002;
395, Padman et al., 1997;
396, Siebenmorgen & Krügel, 2000;
397, Yu et al., 2003b;
398, Clark, 1991;
399, Gueth et al., 1997;
400, Bally et al, 1995;
401, Rodríguez & Reipurth, 1996;
402, Momose et al., 1998;
403, Hilton & Lahulla, 1995;
404, Chini et al., 1997;
405, McMullin et al., 1994;
406, Hurt & Barsony, 1996;
407, McCaughrean et al., 1994;
408, Gibb & Davis, 1998;
409, Bontemps et al., 1995;
410, Bachiller et al., 1995b;
411, Cernicharo & Reipurth, 1996;
412, Nagar et al., 1997;
413, Devine et al., 1999;
414, Codella et al., 2001;
415, Narayanan et al., 2002;
416, Zinchenko et al., 1995.
417, Braz & Scalise, 1982;
418, Furuya et al., 2003;
419, Sun & Wu, 2003;
420, Zhu et al., 2003;
421, Qin & Wu, 2003b;
422, Visser et al., 2001;
423, Visser et al., 2002;
424, Wu & Evans, 2003.

References

- Altenhoff, W. J., Downes, D., Pauls, T. & Schraml, J., 1978, A&AS 35, 23
- André, Ph., Martín-Pintado, J., Despois, D. & Montmerle, T., 1990, A&A 236, 180
- André, P., Motte, F. & Bacmann, 1999, ApJ 513, L57
- Anglada, G. & Rodríguez L. F., 2002, *Revista Mexicana de Astronomía y Astrofísica* 38, 13
- Arce, H. G. & Goodman, A. A., 2001, ApJ 554, 132
- Arce, H. & Goodman, A., 2002, ApJ 575, 928
- Armstrong, J. T. 1989, In: Winnewisser, G. & Armstrong, J. T.(eds.), Symp. at Zermatt, *The Physics and Chemistry of Interstellar Molecular Clouds*, Springer-Verlag, p.143
- Armstrong, J. T. & Winnewisser, G., 1989, A&A 210, 373
- Arquilla, R. & Goldsmith, P. F., 1984, ApJ 279, 664
- Arquilla, R. & Kwok, S., 1987, A&A 173, 271
- Aso, Y., Tatematsu, K., Sekimoto, Y., Nakano, T. & Umemoto, T. et al., 2000, ApJS 131, 465
- Aspin, C., 1992, A&A 266, 416
- Aspin, C., 1998, A&A, 335, 1040
- Aspin, C., McCaughrean, M. J., Casali, M. M. & Geballe, T. R., 1991, A&A 252, 299
- Aspin, C. & Reipurth, B., 2000, MNRAS 311, 522
- Avery, L. W., Hayashi, S. S. & White, G. J., 1990, ApJ 357, 524
- Ayala, S., Curiel, S., Raga, A. L., Noriega-Crespo, A. & Salas, L., 1998, A&A 332, 1055
- Bachiller, R. & Cernicharo, J., 1990, A&A 239, 276
- Bachiller, R., Cernicharo, J., Martín-Pintado, J., Tafalla, M. & Lazareff, B., 1990b, A&A 231, 174
- Bachiller, R., Fuente, A. & Tafala, M., 1995a, ApJ 445, L51
- Bachiller, R., Guilloteau, S., Dutrey, A., Planesas, P. & Martín-Pintado, J., 1995b, A&A 299, 857
- Bachiller, R., Martín-Pintado, J. & Planesas, P., 1991, A&A 251, 639
- Bachiller, R., Menten, K. M. & del Rio-Alvarez, S., 1990a, A&A. 236, 461
- Bachiller, R., Terebey, S., Jarrett, T., Martín-Pintado, J. & Beichman, C. A. et al., 1994, ApJ 437, 296
- Bally, J., 1982, ApJ 261, 558
- Bally, J., Castets, A. & Duvert, G., 1994, ApJ 423,310
- Bally, J., Devine, D. & Alten, V., 1996b, ApJ 473, 921
- Bally, J., Devine, D., Fesen, R. A. & Lane, A. P., 1995, ApJ 454, 345
- Bally, J., Devine, D. & Reipurth, B., 1996a, ApJ 473, L49
- Bally, J. & Lada, C. J., 1983, ApJ 265, 824
- Bally, J., Reipurth, B., Walawender, J. & Armond, T., 2002a, AJ 124, 2152
- Bally, J., Reipurth, B., Lada, C.J. & Billawala, Y., 1999, AJ 117, 410
- Barsony, M., 1989, ApJ 345, 268
- Beichman, C. A., Jennings, R. E., Emerson, J. P., Harris, S. & Rowan-Robinson, M. et al., 1984, ApJ 278, L45
- Beichman, C. A., Myers, P. C., Emerson, J. P., Harris, S. & Mathieu, R. et al., 1986, ApJ 307, 337
- Belloche, A., André, P., Despois, D. & Blinder, S., 2002, A&A 393, 927
- Beltrán, M., Girart, J. M., Estalella, R., Ho, P. T. P. & Palau, A., 2002, ApJ 573, 246
- Bence, S. J., Padman, R., Lsaak, K. G., Wiedner, M. C. & Wright, G. S., 1998, MNRAS 299, 965
- Bence, S. J., Richer, J. S. & Padman, R., 1996, MNRAS 279, 866
- Benson, P. J., Myers, P. C. & Wright, E. L., 1984, ApJ 279, L27
- Beuther, H., Schilke, P., Sridharan, T. K., Menten, K. M. & Walmsley, C. M. et al., 2002a, A&A 383, 892
- Beuther, H., Walsh, A., Schilke, P., Sridharan, T. K. & Menten, K. M. et al., 2002b, A&A 390, 289
- Beuther, H., Schilke, P., Gueth, F., McCaughrean, M. & Andersen, M. et al., 2002c, A&A 387, 931
- Blake, G. A., Sandell, G., van Dishoeck, E. F., Groesbeck, T. D. & Mundy L. G. et al. 1995, ApJ 441, 689
- Blitz, L. & Thaddeus, P., 1980, ApJ 241, 676
- Bloomer, J. D., Watson, D. M., Pipher, J. C., Forrest, W. J. & Pipher, J. L. et al., 1998, ApJ 506, 727
- Bontemps, S., André, P., Terebey, S. & Cabrit, S., 1996a, A&A 311, 858
- Bontemps, S., André, P. & Ward-Thompson, D., 1995, A&A 297, 98
- Bontemps, S., Ward-Thompson, D. & André, P., 1996b, A&A 314, 477
- Bourke, T., 1995, In: Latter, W. B., Radford, S. J. E., Jewell, P. R., Mangum, J. G. & Bally, J., *IAU Symposium 170 at Tucson, CO: Twenty-five years of millimeter-wave spectroscopy*, Kulwer Academic Publishers, p.406
- Bourke, T., 2001, ApJ 554, L91
- Bourke, T., L., Garay, G., Lehtineu, K., Köhnenkamp, I. & Launhardt, R., 1997, ApJ, 476, 781
- Brackmann, E. & Scoville, N., 1980, ApJ 242, 112
- Brand, J., Cesaroni, R., Caselli, P., Catarzi, M. & Codella, C., et al., 1994, A&AS 103, 541
- Braz, M. A. & Epchtein, N., 1983 A&AS 54, 167

- Braz, M. A. & Scalise, E. Jr., 1982, A&A 107, 272
- Cabrit, S. & Bertout, C., 1992, A&A 261, 274
- Cabrit, S., Goldsmith, P. F. & Snell, R. L., 1988, ApJ 334, 196
- Cabrit, S., Lagage, P. O., McCaughrean, M. J. & Olofsson, G., 1997, A&A 321, 523
- Calvet, N., Cantó, J. & Rodríguez, L. F., 1983, ApJ 268, 739
- Campbell, B. & Thompson, R. I., 1984, ApJ 279, 650
- Cantó, J., Rodríguez, L. F., Barral, J. F. & Carral, P., 1981, ApJ 244, 102
- Cantó, J., Rodríguez, L. F., Calvet, N. & Levreault, R. M., 1984, ApJ 282, 631
- Casali, M. M., Eiroa, C. & Duncan, W. D., 1993, A&A 275, 195
- Casoli, F., Dupraz, C., Cerin, M., Combes, F. & Boulanger, F., 1986, A&A 169, 281
- Cernicharo, J. & Reipurth, B., 1996, ApJ 460, L57
- Cesaroni, R., Felli, M., Testi, L., Walmsley, C. M. & Olmi, L., 1997, A&A 325, 725
- Chakraborty, A., Ojha, D. K., Anandarao, B. G. & Rengarajan, T. N., 2000, A&A 364, 683
- Chandler, C. J. & Carlstrom, J. E., 1996, ApJ 466, 338
- Chandler, C., J., Terebey, S., Barsony, M., Moore, T. J. T. & Gautier, T. N., 1996, ApJ, 471, 308
- Chen, H., Tafalla, M., Greene, T. P., Myers, P. C. & Wilner, D. J., 1997, ApJ, 475, 163
- Chen, Y. F., Yao, Y. Q., Yang, J., Hirao, T. & Ishii, M., 1999, AJ 117, 446
- Chernin, L. M., 1996, ApJ 460, 711
- Chernin, L. M. & Masson, C. R., 1991, ApJ 382, L93
- Chernin, L. M. & Masson, C. R., 1992, ApJ 396, L35
- Chernin, L. M. & Masson, C. R., 1995, ApJ 443, 181
- Chernin, L. M. & Wright, M. C. H., 1996, ApJ 467, 676
- Chini, R., Reipurth, B., Ward-Thompson, D., Bally, J. & Nyman, L.-Å et al., 1997, ApJ 474, L135
- Choi, M., Evans, N. J. II & Jaffe, D. T., 1993, ApJ 417, 624
- Choi, M., Panis, J.-F. & Evans, N. J. II, 1999, ApJS 122, 519
- Churchwell, 1999, In: *The Origin of Stars and Planetary Systems*, Eds: Lada, C. J. & Kylafis, N. D., Kluwer Academic Publishers, p.515-552
- Girart, J. M., Curiel, S., Rodríguez, L. F. & Canto, J., 2002, *Revista Mexicana de Astronomía y Astrofísica* 38, 169
- Clark, F. O., 1986, A&A 164, L19
- Clark, F. O., 1991, ApJS 75, 611
- Codella, C. & Bachiller, R., 1999, ApJ, 350, 659
- Codella, C., Bachiller, R., Nisini, B., Saraceno, P. & Testi, L., 2001, A&A 376, 271
- Cohen, M., 1980, AJ 85, 29
- Cohen, M. & Schwartz, R. D., 1987, ApJ 316, 311
- Comoretto, G., Palagi, F., Cesaroni, R., Felli, M. & Bettarini, A., et al., 1990, A&AS 84, 179
- Correia, J. C., Griffin, M. & Saraceno, P., 1997, A&A, 322, L25
- Davidson, J. A. & Jaffe, D. T., 1984, ApJ 277, L13.
- Davis, C. J., Dent, W. R. F., Matthews, H. E., Aspin, C. & Lightfoot, J. F., 1994a, MNRAS 266, 933
- Davis, C. J., Dent, W. R. F., Matthews, H. E., Coulson, I. M. & McCaughrean, M. J., 2000, MNRAS 318, 952
- Davis, C. J. & Eisloffel, J., 1995, A&A 300, 851
- Davis, C. J., Eisloffel, J. & Ray, T. P., 1994b, ApJ 426, L93
- Davis, C. J., Eisloffel, J. & Smith, M. D., 1996, ApJ 463, 246
- Davis, C. J., Matthews, H. E., Ray, T. P., Dent, W. R. F. & Richer, J. S., 1999b, MNRAS 309, 141
- Davis, C. J., Moriarty-Schieven, G., Eisloffel, J., Hoare, M. G. & Ray, T. P., 1998, AJ 115, 1118
- Davis, C. J., Mundt, R., Eisloffel, J. & Ray, T. P., 1995, AJ 110, 766
- Davis, C. J., Ray, T. P. & Eisloffel, J. & Corcoran, D., 1997, A&A 324, 263
- Davis, C. J., Smith, M. D., Eisloffel, E. F., Davies, J. K., 1999a, MNRAS 308, 539
- Davis, C. J., Stern, L., Ray, T. P. & Chrysostomou, A., 2002, A&A 382, 1021
- Dent, W. R. F. & Aspin C., 1992, MNRAS 259, 401
- Dent, W. R. F., Little, L. T., Kaifu, N., Ohishi, M. & Suzuki, S., 1985a, A&A 146, 375
- Dent, W. R. F., Little, L. T., Sato, S., Ohishi, M. & Yamashita, T., 1985b, MNRAS 217, 217
- Dent, W. R. F., Little, L. T. & White, G. J., 1984, MNRAS 210, 173
- Devine, D., Bally, J., Reipurth B., Shepherd, D. & Watson, A., 1999, ApJ 117, 2919
- Dickel, J. R., Dickel H. R. & Wilson, W. J., 1978, ApJ 223, 840
- Dobashi, K., Onishi, T., Iwata, T., Nagahama, T. & Patel, N., 1993, AJ 105, 1487
- Dobashi, K., Nozawa, S., Hayashi, Y., Sato, F. & Fukui, Y., 1994, AJ 107, 2148
- Dobashi, K., Sato, F. & Mizuno, A., 1998a, PASJ, 50, L15
- Dobash, K. & Uehara, H., 2001, PASJ 53, 799
- Dobashi, K., Yonekura, Y. & Hayashi, Y., 1999, PASJ 53, 811

- Dobashi, K., Yonekura, Y., Hayashi, Y., Sato, F. & Ogawa, H., 1998b, *AJ*, 115, 777
- Duvert, G., Cernicharo, J., Bachiller, R. & Gómez-González, J., 1990, *A&A* 233, 190
- Edwards, S. & Snell, R. L., 1982, *ApJ* 261, 151
- Edwards, S. & Snell, R. L., 1983, *ApJ* 270, 605
- Edwards, S. & Snell, R. L., 1984, *ApJ* 281, 237
- Eiroa, C., Torrelles, J. M., Gómez, J. F., Sakamoto, S. & Hasegawa, T. et al., 1992, *PASJ* 44, 155
- Eisloffel, J., 2000, *A&A* 354, 236
- Eisloffel, J., Smith, M. D. & Davis, C. J., 2000, *A&A* 359, 1147
- Erickson, N. R., Goldsmith, P. F., Snell, R. L., Berson, R. L. & Huguenin, G. R. et al., 1982, *ApJ* 261, L103
- Evans, N. J., II, Balkum, S., Levreault, R. M., Hartmann, L. & Kenyon, S., 1994, *ApJ* 424, 793
- Evans, N. J., II, Beichman, C., Gatley, I., Harvey, P. & Nadeau, D. et al., 1981a, *ApJ* 246, 409
- Evans, N. J., II, Blair, G. N., Harvey, P., Israel, F. & Peters, W. L., III et al., 1981b, *ApJ* 250, 200
- Feunte, A., Neri, R., Martin-Pintado, J., Bachiller, R., Rodríguez-Franco, A. et al., 2002, *A&A* 366, 873
- Fich, M. & Lada, C. J., 1998, *ApJS* 117, 147
- Fich, M. & Lada, C. J., 1997, *ApJ* 484, L63
- Field, D., Lemaire, J. L., Pineau des Forêts, G., Gerin, M. & Leach, S. et al., 1998, *A&A* 333, 280
- Fischer, J., Joyce, R. R., Simon, M. & Simon, T., 1982, *ApJ* 258, 165
- Fischer, J., Sanders, D. B., Simon, M. & Solomon, P. M., 1985, *ApJ* 293, 508
- Frerking, M. A. & Langer, W. D., 1982, *ApJ* 256, 523
- Fukui, Y., 1989, In: Reipurth, B.(ed.) *ESO Workshop on Low Mass Star Formation and Pre-Main Sequence Objects*, p. 95
- Fukui, Y., Iwata, T., Takaba, H., Mizuno, A. & Ogawa, H. et al., 1989, *Nat* 342, 161
- Fukui, Y., Sugitani, K., Takaba, H., Iwata, T. & Mizuno, A. et al., 1986, *ApJ* 311, L85
- Fukui, Y., Takaba, H., Iwata, T. & Mizuno, A., 1988, *ApJ* 325, L13
- Furuya, R. S., Kitamura, Y., Wootten, A., Claussen, M. J. & Kawabe, R., 2003, *ApJS* 144, 71
- Garden, R. P., Geballe, T. R., Gatley, I. & Nadeau, D., 1991b, *ApJ* 366, 474
- Garden, R. P., Hayashi, M., Gatley, I., Hasegawa, T. & Kaifu, N., 1991a, *ApJ* 374, 540
- Garden, R. P., Russel, A. P. G. & Burton, M. G., 1990, *ApJ* 354, 232
- Gibb, A. G. & Little, L. T., 2000, *MNRAS* 313, 663
- Gibb, A. G. & Davis, C. J., 1998, *MNRAS* 298, 644
- Girart, J. M., Curiel, S., Rodríguez, L. F. & Cantó, J., 2002, *Revista Mexicana de Astronomía y Astrofísica* 38, 169
- Goldsmith, P. F., Langer, W. D. & Wilson, R. W., 1986, *ApJ* 303, L11
- Goldsmith, P. F., Snell, R. L., Hemeon-Heyer, M. & Langer, W. D., 1984, *ApJ* 286, 599
- Gomez, M., Whitney, B. A. & Kenyon, S. J., 1997, *AJ* 114, 1138
- Gregersen, E. M., Evans, N. J., II, Mardones, D. & Myers, P. C., 2000, *ApJ* 533, 440
- Gregersen, E. M., Evans, N. J., II, Zhou, S. & Choi, M., 1997, *ApJ* 484, 259
- Gregory, P. C. & Condon, J. J., 1991, *ApJS* 75, 1011
- Gregório Hetem, J. C., Lépine, J. R. D. & Ortiz, R. P., 1990, *Rev. Mexicana Astron. Astrof.* 21, 486
- Griffith, M. R., Wright, A. E., Bruke, B. F. & Ekers, R. D., 1994, *ApJS* 90, 179
- Gueth, F. & Guilloteau, S., 1999, *A&A* 343, 571
- Gueth, F., Guilloteau, S. & Bachiller, R., 1996, *A&A* 307, 891
- Gueth, F., Guilloteau, S., Dutrey, A. & Bachiller, B., 1997, *A&A* 323, 943
- Gueth, F., Schilke, P. & McCaughrean, M. J., 2001, *A&A* 375, 1018
- Haikala, L. & Dietrich, D., 1989, In: Winniewisser, G., Armstrong, J.T. (eds.) *Symp. at Zermatt. The Physics and Chemistry of Interstellar Molecular Clouds*, Springer-Verlag. p.236.
- Haikala, L. K. & Laureijs, R. J., 1989, *A&A* 223, 287
- Hartigan, P., Curiel, S. & Raymond, J., 1989, *ApJ* 347, L31
- Harvey, P. M. & Forveille, T., 1988, *A&A* 197, L19
- Harvey, P. M. & Wilking, B. A., Joy, M., 1984, *ApJ* 278, 156
- Haschick, A. D., Reid, M. J., Burke, B. F., Moran, J. M. & Miller, G., 1981, *ApJ* 244, 76
- Hasegawa, T. I., Mitchell, G. F., Matthews, H. E., & Tacconi, L., 1994, *ApJ* 426, 215
- Hartigan, P., Carpenter, J., Dougados, C. & Skrutskie, M., 1995, *AAS* 187, 2110
- Hayashi, M., Hasegawa, T., Omodaka, T., Hayashi, S. S. & Miyawaki, R., 1987, *ApJ* 312, 327
- Heathcote, S., Morse, J. A., Hartigan, P., Reipurth, B. & Schwartz, R. D. et al., 1996, *AJ* 112, 1141
- Henning, Th., Lapinov, A., Schreyer, K., Stecklum, B. & Zinchenko, I., 2000b, *A&A* 364, 613
- Henning, Th., Schreyer, K., Launhardt, R. & Burkert, A., 2000a, *A&A* 353, 211
- Heyer, M. H., Snell, R. L., Goldsmith, P. F. & Myers, P. C., 1987, *ApJ* 321, 370
- Heyer, M., H., Snell, R. L., Morgan, J. & Schloerb, F. P., 1989, *ApJ* 346, 220
- Hilton, J. & Lahulla, J. F., 1993, *AJ* 106, 672

Hilton, J. & Lahulla, J. F., 1995, A&AS 113, 325

Hilton, J., White, G. J., Cronin, N. J. & Rainey, R., 1986, A&A 154, 274

Hirano, N., Hayashi, S. S., Umemoto, T. & Ukita, N., 1998 ApJ 504, 334

Hirano, N., Kameya, O., Mikami, H., Saito, S. & Umemoto, T. et al., 1997, ApJ 478, 631

Ho, P. T. P., Moran, J. M. & Rodriguez, L. F., 1982, ApJ 262, 619

Hodapp, K.-W. & Ladd, E. F., 1995, ApJ 453, 715

Hofner, P., Kurtz, S., Churchwell, E., Walmsley, C. M. & Cesaroni, R., 1996, ApJ 359, 371

Hogerheijde, M. R., van Dishoeck, E. F., Blake, G. A. & van Langevelde, H. J., 1998, ApJ 502, 315

Holbrook, J., 1996, AAS 28, 1415

Hunter, T. R., Testi, L., Taylor, G. B., Tofani, G. & Felli, M. et al., 1995, A&A 302, 249

Hunter, T. R., Phillips, T. G. & Menten, K. M., 1997, ApJ 478, 283

Hunter, T. R., Taylor, G. B., Felli, M. & Tofani, G., 1994, A&A 284, 215

Hurt, R. L. & Barsony, M., 1996, ApJ 460, L45

Iwata, T., Fukui, Y. & Ogawa, H., 1988, ApJ 325, 372

Jiang, L., Wu, Y. & Miller, M., 2001, Chinese Science Bulletin, in press

Kamazaki, T., Saito, M., Hirano, N., Umemoto, T. & Kawabe, R., 2003, ApJ 584, 357

Kameya, O., Hasegawa, T. I., Hirano, N. & Takakubo, K., 1989, ApJ 339, 222

Kastner, J. H., Weintraub, D. A., Snell, R. L., Sandell, G. & Aspin, C. et al. 1994, ApJ 425, 695

Kawamura, J. H., Hunter, T. R., Tong, C.-Y. E., Blundell, R. & Zhang, Q. et al., 1999, PASP 111, 1088

Kelly, M. L. & Macdonald, G. H., 1996, MNRAS 282, 401

Kenyon, S. J., Hartmann, L., Gopez, M., Carr, J. S. & Tokunaga, A., 1993, AJ 105, 1505

Khanzadyan, T., Smith, M. D., Gredel, R., Stante, T. & Davis, C. J., 2002, A&A 383, 502

Knee, L. B. G., 1992, A&A 259, 283

Knee, L. B. G. & Sandell, G., 2000, A&A 361, 671

Koo, B. C., 1989, ApJ 337, 318

Koo, B. C., Lee, Y., Fuller, G. A., Lee, M. G. & Kwon, S.-M. et al., 1994, ApJ 429, 233

Kumar, M. S., Bachiller, R. & Davis, C. J., 2002, ApJ 576, 313

Kurtz, S., Churchwell, E. & Wood, D. O. S., 1994, ApJS 91, 659

Kutner, M. L., Leung, C. M., Machnik, D. E. & Mead, K. N., 1982, ApJ 259, L35

Kwan, J. & Scoville, N., 1976, ApJ 210, L39

Lada, C. J., 1985, ARA&A 23, 267

Lada, C. J. & Fich, M., 1996, ApJ 459, 638

Lada, C. J. & Gautier, T. N. III, 1982, ApJ 261, 161

Lada, C. J. & Harvey, P. M. 1981, ApJ 245, 58

Lada, C. J., Thronson, H. A. J., Smith, H. A., Schwartz, P. R. & Glaccum, W., 1984, ApJ 286, 302

Ladd, E. F., Adams, F. C., Casey, S., Davidson, J. A. & Fuller, G. A. et al., 1991, ApJ 366, 203

Ladd, E. F. & Hodapp, K.-W., 1997, ApJ 475, 749

Langer, W. D., Castets, A. & Lefloch, B., 1996, ApJ 471, L111

Langer, W. D., Velusamy, T. & Xie, T., 1996, ApJ 468, L41

Lefloch, B., Eislöf, B. & Lazareff, B., 1996, A&A 313, L17

Lee, C., Mundy, L. G., Reipurth, B., Ostriker, E. C. & Stone, J. M., 2000, ApJ 542, 925

Lee, C., Mundy, L. G., Stone, J. M. & Ostriker, E. C., 2002, ApJ 576, 294

Levreault, R. M., 1983, ApJ 265, 855

Levreault, R. M., 1984, ApJ 277, 634

Levreault, R. M., 1985, *Molecular Outflows and Mass Loss in Pre-Main Sequence Stars, Technical Report No.85-1*, Astronomy Dept.Univ. of Texas

Levreault, R. M., 1988, ApJS, 67, 283

Lichten, S. M., 1982a, ApJ 255, L119

Lichten, S. M., 1982b, ApJ 253, 593

Liljestrom, T., Mattila, K. & Friberg, P., 1989, A&A 210, 337

Liseau, R., Sandell, G. & Knee, L. B. G., 1988, A&A 192, 153

Little, L. T., Bergman, P., Cunningham, C. J., Heaton, B. D. & Knee, L. B. G. et al., 1988, A&A 205, 129

Longmore, A. J., Robson, E. I. & Jameson, R. F., 1986, MNRAS 221, 589

Loren, R. B., 1989, ApJ 338, 925

Loren, R. B., Wootten, A. & Wilking, B. A., 1990, ApJ 365, 269

Luhman, M. L., Jaffe, D. T., Sternberg, A., Herrmann, F. & Poglitsch, A., 1997, ApJ 482, 298

Lyder, A., D., Belton, D. S. & Gower, A. C., 1998, AJ 116, 840

Magnier, E. A., Volp, A. W., Laan, K., van den Ancker, M. E. & Waters, L. B. F. M., 1999, A&A 352, 228

Magnier, E. A., Waters, L. B. F. M., Kuan, Y. -J., Chu, Y. -H. & Taylor, A. R. et al., 1995, A&A 305, 936M

Mao, R., 1999, Ph.D thesis, PMO

- Mao, R. Q., Yang, J., Henkel, C. & Jiang, Z. B., 2002, A&A 389, 589
- Mao, X., Wu, Y., Hao, J. & Hou, M., 1989, Universitatis Pekinensis 25, 605
- Mardones, D., Myers, P. C., Tafalla, M. & Wilner, D. J., 1997, ApJ 489, 719
- Margulis, M. & Lada, C. J., 1986, ApJ 309, L87
- Margulis, M., Lada, C. J., Hasegawa, T., Hayashi, S. S. & Hayashi, M. et al., 1990, ApJ 352, 615
- Margulis, M., Lada, C. J. & Snell, R. L., 1988, ApJ 333, 316
- Margulis, M., Lada, C. J. & Young, E. T., 1989, ApJ 345, 906
- Margulis, M. & Snell, R. L., 1989, ApJ 343, 779
- Massi, F., Lorenzetti, D. & Vitali, F., 1997, IAUS 182, 21
- Masson, C. R., Mundy, L. G. & Keene, J., 1990, ApJ 357, L25
- Mathieu, R. D., Benson, P. J., Fuller, G. A., Myers, P. C. & Schild, R. E., 1988, ApJ 330, 385
- Mattila, K., Liljeström, T. & Toriseva, M., 1989, In: Reipurth, B. (ed.) *ESO Workshop on Low Mass Star Formation and Pre-Main Sequence Objects*. p.153
- McCutcheon, W. H., Dewdney, P. E., Purton, C. R. & Sato, T., 1991, AJ 101, 1435
- McMcaughrean, M. J., Rayner, J. T. & Zinnecker, H., 1994, ApJ 436, L189
- McMullin, J. P., Mundy, L. G., Wilking, B. A., Hezel, T. & Blake, G. A., 1994, ApJ 424, 222
- Mendoza, E. E., Rodríguez, L. F., Chavarria-K, C. & Neri, L., 1990, MNRAS 246, 518
- Meyers-Rice, B. A. & Lada, C. J., 1991, ApJ 368, 445
- Minchin, N. R., Sandell, G. & Murray, A. G., 1995, A&A 293, L61
- Minchin, N. R., White, G. J. & Padman, R., 1993, A&A 277, 595
- Mitchell, G. F. & Hasegawa, T. I., 1991, ApJ 371, L33
- Mitchell, G. F., Lee, S. W., Maillard, J.-P., Matthews, H. & Hasegawa, T. I. et al., 1995, ApJ 438, 794
- Mitchell, G. F., Maillard, J.-P. & Hasegawa, T. I., 1991, ApJ 371, 342
- Mizuno, A., Fukui, Y., Iwata, T., Nozawa, S. & Takano, T., 1990, ApJ 356, 184
- Molinari, S., Testi, L., Brand, J., Cesaroni, R. & Palla, F., 1998, ApJ 505, L39
- Momose, M., Ohashi, N., Kawabe, R., Nakano, T. & Hayashi, M., 1998, ApJ 504, 314
- Monin, J.-L., Pudritz, R. E., & Lazareff, B., 1996, A&A 305, 572
- Moran, J. M. & Rodríguez, 1980, ApJ 236, L159
- Moreira, M., Jessop, N., Santos, C. & Yun, J., 2000, AJ 119, 2960
- Morgan, J. A. & Bally, J., 1991, ApJ 372, 505
- Morgan, J. A., Schloerb, F. P., Snell, R. L. & Bally, J., 1991, ApJ 376, 618
- Moriarty-Schieven, G. H., Butner, H. M. & Wannier, P. G., 1995, ApJ 445, L55
- Moriarty-Schieven, G. H. & Snell, R. L., 1988, ApJ 332, 364
- Moriarty-Schieven, G. H. & Wannier, P. G., 1991, ApJ 373, L23
- Moriarty-Schieven, G. H., Wannier, P. G., Tamura, M. & Keene, J., 1992, ApJ 400, 260
- Moro-Martín, A., Cernicharo, J., Noriega-Crespo, A. & Martín-Pintado, J., 1999, ApJ 520, L111
- Mozurkewich, D., Schwartz, P. R. & Smith, H. A., 1986, ApJ 311, 371
- Myers, P. C., Bachiller, R., Caselli, P., Fuller, G. A. & Mardones, D. et al., 1995, ApJ 449, L65
- Myers, P. C., Fuller, G. A., Mathieu, R. D., Beichman, C. A. & Benson, P. J. et al., 1987, ApJ 319, 340
- Myers, P. C., Heyer, M., Snell, R. L. & Goldsmith, P. F., 1988, ApJ 324, 907
- Myers, P. C., Mardones, D., Tafalla, M., William, J. P. & Wilner, D. J., 1996, ApJ 465, L133
- Nagar, N. M., Vogel, S. N., Stone, J. M. & Ostriker, E. C., 1997, ApJ
- Nakano, M. & Yoshida, S., 1986, PASJ 38, 531
- Narayanan, G., Moriarty-Schieven, G., Walker, C. K. & Butner, H. M., 2002, ApJ 565, 319
- Nirano, N., Kamega, O., Mikami, H., Saito, S. & Umemoto, T. et al., 1997, ApJ, 478, 631
- Nisini, B., Massi, F., Vitali, F., Giannini, T. & Lorenzetti, D. et al., 2001, A&A 376, 553
- Noriega-Crespo, A., Cotera, A., Young, E. & Chen, H., 2002, ApJ 580, 959
- Odenwald, S. F. & Schwartz, P. R., 1993, ApJ 405, 706
- Ogura, K., Sugitani, K. & Pickels, A., 2002, AJ 123, 2597
- Ohashi, N., Hayashi, M., Ho, P. T. P., Momose, M. & Tamura, M. et al., 1997, ApJ 488, 317
- Olberg, M., Reipurth, B. & Booth, R. S., 1992, A&A 259, 252
- Olberg, M., Reipurth, B. & Booth, R. S., 1989, In: Winnewisser, G., Armstrong, J. T. (eds.) *Sump. at Zermatt. The Physics and Chemistry of Interstellar Molecular Clouds*. Springer-Verlag, p.120
- Olmi, L., Felli, M. & Cesaroni, R., 1997, A&A 326, 373
- Olofsson, H., Ellder, J., Hjalmarsen, A. & Rydbeck, G., 1982, A&A 113, L18
- Onishi, T., Kawamura, A., Abe, R., Yamaguchi, N. & Saito, H. et al., 1999, PASJ 51, 871
- Padman, R., Bence, S. & Richer, J., 1997, IAU 182, 123
- Palacios, J., & Eiroa, C., 1999, A&A 346, 233
- Parker, N. D., Padman, R., Scott, P. F. & Hills, R. E., 1988, MNRAS 234, 67P
- Parker, N. D., Padman, R. & Scott, P. F., 1991, MNRAS 252, 442

- Patel, N. A., Greenhill, L. J., Herrnstein, J., Zhang, Q. & Moran, J. M. et al., 2000, ApJ 538, 268
- Phillips, J. P. & Mampaso, A., 1991, A&AS 88, 189
- Phillips, J. P. & Mampaso, A., 1996, A&A 316, 182
- Phillips, J. P., White, G. J., Rainey, R., Avery, L. W. & Richardson, K. J. et al., 1988, A&A 190, 289
- Poetzel, R., Mundt, R. & Ray T. P., 1992, A&A 262, 229
- Porrás, A., Curz-González, I. & Salas, L., 2000, A&A 361, 660
- Qin, S. L. & Wu, Y. F., 2003b, *Acta Astronomica Sinica* Suppl. 44, 89
- Qin, S. & Wu, Y., 2003a, CJA&A 3, 69
- Reipurth, B., 1999, *A General Catalogue of HH Objects*, Electronic Version
- Reipurth, B., 1989, In: Reipurth, B.(ed.) *ESO Workshop on Low Mass Star Formation and Pre-Main Sequence Objects*, p.247
- Reipurth, B., Chini, R., Krügel, E., Kreysa, E. & Sievers, A., 1993, A&A 273, 221
- Reipurth, B. & Graham, J. A., 1991, In: Reipurth, B.(ed.) *Low Mass Star Formation in Southern Molecular Clouds*, ESO Scientific Report 11, 149
- Reipurth, B., Heathcote, S., Yu, K. C., Bally, J. & Rodríguez L. F., 2000, ApJ 534, 317
- Reipurth, B. & Olberg, M., 1991, A&A 246, 535
- Reipurth, B., Olberg, M., Gredel, R. & Booth, R., S., 1997, A&A, 327, 1164
- Richer, J. S., 1990, MNRAS 245, 24p
- Richer, J. S., Hills, R. E. & Padman, R., 1992, MNRAS 254, 525
- Richer, J. S., Hills, R. E., Padman, R. & Russell, A. P. G., 1989 MNRAS 241, 231
- Ridge, N. A. & Moore, T. J. T., 2001, A&A 378, 495
- Rodríguez, L. F. & Canto, J., 1982, ApJ 255, 103
- Rodríguez, L. F., Carral, P., Ho, P. T., P. & Moran, J. M., 1982, ApJ 260, 635
- Rodríguez, L. F., Ho, P. T. P. & Moran, J. M., 1980, ApJ 240, L149
- Rodríguez, L. F. & Reipurth, B., 1996, *Revista Mexicana de Astronomía y Astrofísica* 32, 27
- Rodríguez, L. F., Torrelles, J. M. & Moran, J. M., 1981, AJ 86, 1245
- Ryder, S. D., Allen, L. E., Burton, M. G., Ashley, M. C. B. & Storey, J. W. V., 1998, MNRAS 294, 338
- Saito, H., Tachihara, K., Onishi, T., Yamaguchi, N. & Mizuno N. et al., 1999, PASJ 51, 819
- Sandell, G., Avery, L. W., Bass, F., Coulson, I. & Dent, W. R. F. et al., 1999, ApJ 519, 236
- Sandell, G. & Aspin, C, 1998, A&A 333, 1016
- Sanders, D. B. & Willner, S. P., 1985, ApJ 293, L39
- Sato, F. & Fukui, Y., 1989, ApJ 343, 773
- Sato, F., Mizuno, A., Nagahama, T., Onishi, T. & Yonekura, Y. et al., 1994, ApJ 435, 279
- Scalise, E. Jr., Rodriguez, L. F. & Mendoza-Torres, E., 1989, A&A 221, 105
- Schmid-Burgk, J., Densing, R., Krugel, E., Nett, H. & Roser, H. P. et al., 1989, A&A 215, 150
- Schmid-Burgk, J., Güsten, R., Mauersberger, R., Schulz, A. & Wilson, T. L., 1990, ApJ 362, L25.
- Schreyer, K., Helmich, F. P., van Dishoeck, E. F. & Henning, Th., 1997, A&A 347, 365
- Schwartz, P. R., Frerking, M. A. & Smith, H. A., 1985, ApJ 295, 89
- Schwartz, P. R., Gee, G. & Huang, Y. L., 1988, ApJ 327, 350
- Schwartz, P. R., Thronson, H. A. Jr, Lada, C. J., Smith, H. A. & Glaccum, W. et al., 1983, ApJ 271, 625
- Scoville, N. Z., Sargent, A. I., Sanders, D. B., Claussen, M. J. & Masson, C. R. et al., 1986, ApJ 303, 416
- Shepherd, D. S. & Churchwell, E., 1996a, ApJ 457, 267
- Shepherd, D. S. & Churchwell, E., 1996b, ApJ 472, 225
- Shepherd, D. S. & Churchwell, E. & D. J. Wilner, 1997, ApJ 482, 355
- Shepherd, D. S., Yu, K. c., Bally, J. & Testi, L., 2000, ApJ 535, in press
- Shepherd, D. S. & Kurtz, S., 1998, AAS 193, 7301s
- Shepherd, D. S., Watson, A. M., Sargent, A. J. & Churchwell, E., 1998, ApJ 507, 861
- Shi, Y., Wu, Y., Miller, M. & Mao, R. Q., 2003, *Acta Astronomica Sinica* Suppl. 44, 92
- Siebenmorgen, R. & Krügel, E., 2000, A&A 364, 625
- Simon, M., Cassar, L., Felli, M., Fischer, J. & Massi, M. et al., 1984. ApJ 278, 170
- Smith, H. A. & Fischer, J., 1992, ApJ 398, L99
- Snell, R. & Bally, J., 1986, ApJ 303, 683
- Snell, R. L., Dickman, R. L. & Huang, Y.-L., 1990, ApJ 352, 139
- Snell, R. L. & Edwards, S., 1981, ApJ 251, 103
- Snell, R. L. & Edwards, S., 1982, ApJ 259, 668
- Snell, R. L., Huang, Y. L., Dickman, R. L. & Claussen, M. J., 1988, ApJ 325, 853
- Snell, R. L., Loren, R. B. & Plambeck, R. L., 1980, ApJ 239, L17
- Snell, R. L., Scoville, N. Z., Sanders, D. B. & Erickson, N. R., 1984, ApJ 284, 176
- Stanke, T., McCaughrean, M. J. & Zinnecker, H., 1998a, A&A 332, 307

- Stanke, T., McCaughrean, M. J. & Zinnecker, H., 1998b, *Astronomische Gesellschaft Meeting Abstracts, Abstracts of Contributed Talks and Posters presented at the Annual Scientific Meeting of the Astronomische Gesellschaft at Heidelberg* 14, 16
- Stanke, T., McCaughrean, M. J. & Zinnecker, H., 2000, *A&A* 355, 639
- Stanke, T., McCaughrean, M. J. & Zinnecker, H., 2002, *A&A* 392, 239
- Strom, K. M., Strom, S. E. & Vrba, F. J., 1976, *AJ* 81, 320
- Sugitani, K., Fukui, Y., Mizuno, A. & Ohashi, N., 1989, *ApJ* 342, L87
- Sugitani, K., Fukui, Y. & Ogura, K., 1991, *ApJS* 77, 59
- Sugitani, K. & Ogura, K., 1994, *ApJS* 92, 163
- Sun, F. & Wu, Y., 2003, *Ch.A&A* 27, 73
- Tamura, M., Gatley, I., Waller, W. & Werner, M. W., 1991, *ApJ* 374, L25
- Tamura, M., Ohashi, N., Hirano, N., Itoh, Y. & Moriarty-Schieven, G. H., 1996, *AJ* 112, 2076
- Tamura, M., Sato, S., Suzuki, H., Kaifu, N. & Hough, J. H., 1990, *ApJ* 350, 728
- Terebey, S., Beichman, C. A., Gautier, T. N., & Hester, J. J., 1990, *ApJ* 362, L63
- Terebey, S., Vogel, S. N. & Myers, P. C., 1989, *ApJ* 340, 472
- Torrelles, J. M., Verdes-Montenegro, L., Ho, P. T., Rodríguez, L. F. & Canto, J., 1993, *ApJ* 410, 202
- Umemoto, T., Hirano, N., Kameya, O., Fukui, Y. & Kuno, N. et al., 1991, *ApJ* 377, 510
- Umemoto, T., Iwata, T., Fukui, Y., Mikami, H. & Yamamoto, S. et al., 1992, *ApJ* 392, L83
- Valdettaro, R., Palla, F., Brand, J., Cesaroni, R. & Comoretto, G. et al., 2001, *A&A* 368, 845
- Vallée, J. P. & Avery, L. W., 1990, *A&A* 233, 553
- Velusamy, T. & Langer, W. D., 1998, *Nature* 392, 685
- Visser, A. E., Richer, J. S. & Chandler, C. J., 2001, *MNRAS* 323, 257
- Visser, A. E., Richer, J. S. & Chandler, C. J., 2002, *AJ* 124, 2756
- Walker, C. K., Lada, C. J., Young, E. T., Maloney, P. R. & Wilking, B. A., 1986, *ApJ* 309, L47
- Walker, C. K., Lada, C. J., Young, E. T. & Margulis, M., 1988, *ApJ* 332, 335
- Walsh, A. J., Burton, M. G., Hyland, A. R. & Robinson, G., 1998, *MNRAS* 301, 640
- Wang, Y., Evans, N. J. II, Zhou, S. & Clemens, D. P., 1995, *ApJ* 454, 217.
- Ward-Thompson, D., Buckley, H. D., Greaves, J. S., Holland, W. S. & André, P., 1996, *MNRAS* 281, L53
- Weintraub, D. A., Kastner, J. H., Griffith, L. L. & Campins, H., 1993, *AJ* 105, 271
- White, G. J., 1993, *A&A* 274, L33
- White, G. J., Casali, M. M. & Eiroa, C., 1995, *A&A* 298, 594
- White, G. J. & Phillips, J. P., 1981, *MNRAS* 194, 947
- White, G. J., Tothill, N. F. H., Matthews, H. E., McCutcheon, W. H. & Hultgren, M. et al., 1997, *A&A* 323, 529
- Wilking, B. A., Blackwell, J. H. & Mundy, L. G., 1990, *AJ* 100, 758
- Wilson, T. L. & Mauersberger, R., 1991, *A&A* 244, L33
- Wolf, G. A., Lada, C. J. & Bally, J., 1990, *AJ* 100, 1892
- Wolf-Chase, G. A., Barsony, M. & O'Linger, J., 2000, *AJ* 120, 1467
- Wolf-Chase, G. A. & Gregersen, E., 1997, *ApJ* 479, L67
- Wootten, A., 1989, *ApJ* 337, 858
- Wootten, A. & Loren, R. B., 1987, *ApJ* 317, 220
- Wouterloot, J. G. A. & Brand, J., 1999, *A&AS* 140, 177
- Wouterloot, J. G. A., Brand, J. & Henkel, C., 1988, *A&A* 191, 323
- Wouterloot, J. G. A. & Brand, J., 1989, *A&AS* 80, 149
- Wouterloot, J. G. A., Henkel, C. & Walmsley, C. M., 1989, *A&A* 215, 131
- Wright, A. E., Griffith, M. R., Burke, B. F. & Ekers, R. D., 1994, *ApJS* 91, 111
- Wu, J. & Evans, N. J., II, 2003, *ApJ* 592, L79
- Wu, Y., Yang, C. & Li, Y., 1999b, *CA&A* 23, 6
- Wu, Y., Yang, C., Li, Y., Lei, C., Sun, J., et al., 1999a, *Science in China(Ser. A)*, 42, 732
- !Wu, Y. & Yu, W., 2003, preprint
- Wu, Y., Zhang, Q., Chen, H., Yang, C., Wei, Y. & Ho, P. T. P., 2004, *AJ*, submitted
- Wu, Y., Zhou, S. & Evans, N. J. II, 1992, *ApJ* 394, 196
- Wynn-Williams, C. G., 1986, *ARA&A* 20, 587
- Xiang, D. & Turner, B. E., 1995, *ApJS*, 99, 121
- Xie, T. & Goldsmith, P. F., 1990, *ApJ* 359, 378
- Yamaguchi, N., Mizuno, N., Saito, H., Matsunaga, K. & Mizuno, A. et al., 1999a, *PASJ* 51, 775
- Yamaguchi, N., Mizuno, N., Saito, H., Matsunaga, K. & Mizuno, A. et al., 1999b, *Proceedings of Star Formation 1999*
- Yamashita, T., Sato, S., Nagata, T., Gatley, I. & Hayashi, S. S. et al., 1989b, *ApJ* 336, 832
- Yamashita, T., Suzuki, H., Kaifu, N., Tamura, M. & Mountain, C. M. et al., 1989a, *ApJ* 347, 894

- Yang, C. & Wu, Y., 1998, *CA&A*, 22, 451
- Yang, J., Fukui, Y., Umemoto, T., Ogawa, H. & Chen, H., 1990, *ApJ* 362, 538
- Yang, J., Ohashi, N., Yan, J., Liu, C. & Kaifu, N. et al., 1997, *ApJ* 475, 683
- Yang, J., Umemoto, T., Iwata. T. & Fukui, Y., 1991, *ApJ* 373, 137
- Yonekura, Y., Dobashi., K., Hayashi, Y., Sato, F. & Ogawa, H., 1998, *AJ* 115, 2009
- Yu, K. C., Bally, J. & Hereld, M., 1995, *AAS* 187, 2113
- Yu, K. C., Bally, J. & David, D., 1997, *ApJ* 485, L45
- Yu, K. C., Billawala, Y., Smith, M. D., Bally, J. & Butner, H. M., 2000, *AJ* 120, 1974
- Yu, K. C., Billawala, Y. & Bally, J., 1999, *AJ* 118, 2940
- Yu, W., Wu., Y. & Miller, M., 2003b, preprint
- Yu, W., Wu, Y. & Wang, L., 2003a, *Acta Astronomica Sinica* Suppl. 44, 101
- Yun, J. L. & Clemens, D. P., 1992, *ApJ* 385, L21.
- Yun, J. L. & Clemens, D. P., 1994, *ApJS* 92, 145.
- Yun, J. L., Clemens, D. P., Moreira, M. C. & Santos, N. C., 1997, *ApJ* 479, L71
- Yun, J. L., Santos, C. A., Clemens, D. P., Afonso, J. M. & McCaughrean, M. J. et al., 2001, *A&A* 372, L33
- Zhang, Q., Hunter, T. R., Brand, J., Sridharan & T. K., Molinari et al., 2001, *ApJ* 552, L167 (The data is taken from private communication.)
- Zhao, B., Wang, M., Yang, J., Wang, H. & Deng, L. et al., 1999, *AJ* 118, 1347
- Zhao, M., Wu, Y., Miller, M. & Mao, R., 2003, *Acta Astronomica Sinica* Suppl. 44, 103
- Zhou, S., 1995, *ApJ* 442, 685
- Zhou, S., *Low Mass Star Formation in Millimeter-Wave Astronomy: Molecular Chemistry & Physics in Space*, 199-230, W. F. Wall et al.(eds.), 1999 Kulwer Academic Publishers
- Zhou, S., Evans, N. J. II, Gusten, R., Mundy, L. G. & Kutner, M. L., 1991, *ApJ* 372, 518
- Zhou, S., Evans, N. J. II, Kömpe, C. & Walmsley, C. M., 1993, *ApJ* 404, 232
- Zhou, S., Evans, N. J. II & Wang, Y., 1996, *ApJ* 466, 296
- Zeilik, M. II & Lada, C. J., 1978, *ApJ* 222, 896
- Zinchenko, I., Mattila, K. & Toriseva, M., 1995, *A&AS* 111, 95
- Ziurys, L. M., Wilson, T. L. & Mauetsberger, R., 1990, *ApJ* 356, L25
- Zhu, L., Wu, Y. & Wei, Y., 2003, in preparation
- Zuckerman, B., Kuiper, T. B. H. & Kuiper, E. N. R., 1976, *ApJ* 209, L137

Table 1a CO Molecular Outflows

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
001	LKH $_{\alpha}$	00 08 47	+58 33 06	117 46	−03 38	1.0	−63	13	Bi	0.85	<u>3.4</u>	1E3
002	00117+6412	00 11 45	+64 12 04	118 57	+01 53	3.3		8.6	Bi	0.81		4.6E5
003	00213+6530	00 21 22	+65 30 25	120 08	+03 04	0.85	45	14.4	Bi	1.11	2.3	12.9
004	00259+6510	00 25 58	+65 10 12	120 35	+02 41	0.85	40	17.5	Bi	1.51	3.2	9.9
005	00259+5625 CB3	00 25 59	+56 25 32	119 48	−06 02	0.6	−263	11	Bi	0.33	1.2	830
006	00338+6312 L1287	00 33 53	+63 12 32	121 18	+00 40	1.6	18	9.5	Bi	0.64	2.0	930
	RNO1b/1c							25				2.4E3
007	00342+6374	00 34 18	+63 47 45	121 23	+01 14	2	43	40	Multi	1.19	4.1	
008	00420+5530	00 42 05	+55 30 54	122 01	−07 04	7.72	−950	75	Bi	2.20	1.09	5.15E4
009	00494+5617	00 49 28	+56 17 28	123 04	−06 19	2.2	−242	20	Bi	1.32	1.4	5.4E3
010	NGC281-E	00 51 18	+56 17 07	123 19	−06 19	2.1	−231		Bi			
011	01133+6434	01 13 18	+64 34 50	125 36	+02 06	3.0	110	10.4	Bi	0.76	3.7	3E3
012	W3-IRS5	02 21 53	+61 52 21	133 43	+01 13	2.3	49	52	Bi			11E5
								72				2E5
								47				1.1E6
013	IC1805-W	02 25 15	+61 20 10	134 17	+00 51	2.3	34		Bi			1100
014	02310+6133	02 30 56	+61 33 15	134 49	+01 19	3.8	87		Bi			
015	02461+6147	02 46 12	+61 47 34	136 23	+02 19	4.2	166		Bi			1.6E6
016	AFGL4029	02 57 36	+60 17 22	138 18	+01 33	2.2	60	5.5	Bi	1.49	2.3	2.1E4
	02575+6017											
017	AFGL437	03 03 32	+58 19 37	139 54	+00 12	2.0	7	24	Bi	1.2	<u>1.6</u>	(0.2-1)E4
								23.0				2.40E4
018	RNO13	03 22 05	+30 35 50	157 57	−21 27	0.2	−73	12	MB	0.1	2.5	5.2
019	L1448 IRS2	03 22 17.9	+30 34 41	158 00	−21 26	0.3	−110	24	Bi			
020	L1448 IRS3	03 22 32	+30 34 49	158 03	−21 25	0.3	−110	70	Bi	0.08	~ 2.0	
										0.20		9
021	L1448 U-star	03 22 34	+30 33 34	158 04	−21 25	0.3	−110	140	Bi	0.18	~ 5.4	3
022	AFGL490-iki	03 23 28	+58 43 44	141 55	+01 54	0.975	32	48	Bi	0.77	3.1	1100
	03234+5843											
023	AFGL490	03 23 39	+58 36 39	141 60	+01 49	0.9	29	65	Bi	0.62	<u>1.9</u>	1.4E3
								161				1.4-2.4E3
												1.4E3
												1.7-2.4E3
024	L1455NW	03 24 22	+30 05 24	158 42	−21 34	0.35	−129	10	Bi	0.42	<u>3.7</u>	
												~0.3-0.7
025	RNO15FIR L1445 FIR	03 24 36	+30 02 40	158 46	−21 34	0.35	−129	15	MB			7.2
												1.5
												6.4
026	L1455M	03 24 44	+30 01 43	158 48	−21 34	0.35	−129	16	Bi	0.44	<u>2.8</u>	25
027	NGC1333	03 25 34	+31 03 14	158 18	−20 38	0.50	−176	10.5	Bi	0.32	1.0	
028	IRAS2NNE-SSW	03 25 48	+31 04 34	158 20	−20 35	0.22	−123		Bi	0.26		
029	IRAS2east-west	03 25 50.1	+31 04 18	158 20	−20 36	0.22	−123		Bi	0.237		
030	HH12	03 25 57.5	+31 05 34	158 18	−20 30	0.35	−122		MB	0.15		23
								20				30
031	HH7-11 SSV	03 25 58.1	+31 05 45	158 21	−20 33	0.50	−123		Bi	0.40	<u>1.9</u>	90
												150
								40				
032	NGC1333 IRAS4A	03 26 05.0	+31 03 14	158 24	−20 34	0.22	−123	46	Bi	0.33		
033	HH6	03 26 05.2	+31 08 01	158 21	−20 30	0.22	−123		Bi	0.15		
034	03271+3013	03 27 10	+30 13 34	159 08	−21 06	0.35	−126	80	Bi			≤ 3.0
035	03282+3035	03 28 15	+30 35 14	159 06	−20 40	0.35	−124	140	Bi	1.60	> 5.9	≤ 1.5
												3
036	03301+3057 Barnard 1	03 30 17	+30 57 26	159 13	−20 06	0.35	−120	40	Bi	0.01		2.4
037	HH211	03 40 49	+31 51 24	160 29	−18 00	0.32	−99	60	Bi	0.09	16.8	
038	B5-IRS3	03 43 56	+32 33 54	160 33	−17 02	0.35	−102	10	MR	0.43	2.2	> 0.5
039	B5-IRS1	03 44 32	+32 42 30	160 39	−16 57	0.35	−10	31	Bi	0.3		9.6
											<u>6</u>	
040	B5-IRS4	03 44 36	+32 54 33	169 34	−16 50	0.35		6	Bi			> 0.3

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
041	B5-IRS2	03 44 54	+32 40 17	156 41	-19 45	0.35	-10	10	Bi	1.4		1.3
042	L1489	04 01 41	+26 10 48	168 04	-19 10	0.14	-46	5.5	Bi	0.14	<u>2.0</u>	4.2
043	PP 13S	04 07 21	+38 00 08	160 30	-09 48	0.35	-60	29	Bi			30
044	04166+2706	04 16 38	+27 06 29	169 50	-16 08	0.14	-39	11.5	Bi			0.4
045	04181+2655	04 18 06	+26 55 01	170 12	-16 01	0.14	-39	10	Bi	> 0.056	> 1.57	0.5
046	T Tau	04 19 04	+19 25 05	176 14	-20 53	0.14	-50	13	MB	0.15	<u>1.0</u>	22
						0.14		30				18.3
						0.16						10
047	04191+1523	04 19 09	+15 23 16	179 33	-23 29	0.14	-56		Bi			< 210
	IRAM04191					0.14			Multi			0.15
048	04239+2436	04 23 55	+24 36 54	172 54	-16 36	0.14	-40	24.7	MR			1.1
049	Haro 6-10	04 26 22	+24 26 26	173 25	-16 18	0.14	-39		Bi			5.0
	L1524											
050	ZZ Tau	04 27 51	+24 35 24	173 32	-15 57	0.14	-38	5.2	MR	0.14	<u>2.0</u>	0.18
051	L1551-IRS5	04 28 40	+18 01 42	178 56	-20 03	0.12	-41	30	Bi	0.42	<u>5.2</u>	25
						0.16						33
052	L1551NE	04 28 43	+18 02 12	178 56	-20 02	0.16	-55	18	MR			6
053	HL/XZ Tau	04 28 44	+18 07 36	178 51	-19 58	0.144	-48	11	MB			8.5
						0.16						4.4-10.7
054	TMC2A ^b	04 28 58	+24 24 27	173 51	-15 54	0.14	-38	8.8	Bi			1.4
												1.07
055	L1529	04 29 43	+24 16 54	174 04	-15 50	0.14	-38	30	Bi			2
056	04325+2402	04 32 32	+24 02 08	174 41	-15 30	0.14	-37	9.0	MR	0.93	<u>2.4</u>	0.41
	L1535											0.3
057	L1642	04 32 32	-14 19 18	210 49	-36 36	0.13	-60	3.6	Bi	0.1	3.3	0.3
058	L1527	04 36 49	+25 57 16	173 49	-13 32	0.16	-37	6.0	Bi			1.30
	L1527 IRS											2.8
059	04361+2547	04 36 10	+25 47 30	173 51	-13 45	0.14	-33		Bi	0.013	~ 2.3	1.9
060	TMC1A	04 36 31	+25 35 55	174 03	-13 48	0.14	-33	11	Bi	0.02	2.6	0.86
061	IC2087	04 36 55	+25 39 17	174 04	-13 42	0.14	-33	7.3	MR	0.72	<u>2.1</u>	0.73
062	04381+2540	04 38 09	+25 40 53	174 14	-13 28	0.14	-33	36.6	Bi	0.03	2.8	0.5
	TMC1											5.61E4
063	05137+3919	05 13 46	+39 19 10	168 04	+00 49	10.80	155	55	Bi	3.98	2.7	2.40E4
064	HH114/115	05 15 35	+07 07 54	195 16	-16 59	0.46	-134		Multi			16
065	05168+3634	05 16 54	+36 34 21	170 39	-00 16	6.08	-28	66	Bi	2.73	1.78	8.1E3
066	L1634	05 17 22	-05 55 05	207 36	-23 02	0.5	-196		Bi			3.80E3
067	AFGL5142	05 27 28	+33 45 37	174 12	-00 05	1.8	-2	12	Bi	0.67	1.2	
	05274+3345					1.8		23				
									Multi			
068	L1582B	05 29 27	+12 47 13	192 09	-11 07	0.46	-89	12	Bi	0.13	2.6	5.9
069	RNO43 ¹	05 29 33	+12 47 33	192 09	-11 06	0.40	-88		Bi			12
						0.46						
070	RNO43S	05 29 35	+12 50 00	192 08	-11 03	0.46	-88	11	Bi	0.27	1.4	
071	RNO43N	05 29 38	+12 55 00	192 04	-11 00	0.46	-88	11	Bi	0.40	2.8	
072	Ori A-W	05 30 15	-05 37 52	208 54	-20 03	0.50	-171	13	Bi			40-400
073	HH83	05 31 06	-06 31 45	209 51	-20 16	0.47	-163	9.5	Bi			8
074	Orion-S	05 32 45	-05 26 05	209 01	-19 24	0.48	-166	50	Bi	0.21	4.2	8.5E3
075	CSO 2	05 32 46	-05 01 15	211 00	-14 47	0.45			Bi	0.14		
076	05327+3404	05 32 47	+34 04 20	174 32	+01 01	1.2-1.6	21 - 28	~20	Bi			82
								16.53				1.1E4
077	Orion-A	05 32 47	-05 24 14	208 59	-19 23	0.5	-166	266	Bi	0.04	<u>1.0</u>	1-10E4
	Orion-KL							127				2-10E4
												2.1E5
												2.1E5
078	Orion-KLN	05 32 48	-05 22 34	208 57	-19 22	0.5	-166	28.3	Bi			< 56
079	AC 3	05 32 50.4	-05 02 34	208 39	-19 12	0.45	-148		MB	0.41		128
080	FIR 1	05 32 54	-05 09 30	208 46	-19 15	0.45	-148		Bi	0.17		

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
081	OMC-2/3(MMS 9)	05 32 58	-05 07 30	208 45	-19 13	0.45	-148		Bi			≤ 94
082	OMC-2	05 33 00	-05 11 32	208 49	-19 14	0.5	-165	13	Bi	0.18	1.9	
083	HH 34	05 33 04	-06 28 49	210 25	-19 49	0.46	-155	13	Bi			
084	MMS 10	05 33 06	-05 07 30	208 46	-19 11	0.45			MB	0.31	2.4	
085	L1641N	05 33 53	-06 24 02	210 04	-19 36	0.50	-168	28.0	Bi	0.21	2.0	40-400
						0.45		44.8				$10^{2.1}$
086	CS-star HH1 VLA1	05 33 55	-06 47 24	210 26	-19 46	0.48 0.44	-162	8.0 21 11.05	MR	0.08	5.2	36
087	05339-0647	05 33 56.9	-06 46 46	210 26	-19 45	0.44	-148	12	Bi			
088	VLA3	05 33 57	-06 47 10	210 26	-19 45	0.47	-250	40.0	Bi	0.1	3.3	
089	VLA 2	05 33 57.0	-06 46 46	210 26	-19 45	0.44	-149		MR			
090	HH 1-2	05 33 57	-06 47 55	210 26	-19 46	0.46	-155	13	Bi			
091	05339-0646	05 33 58	-06 46 33	210 26	-19 44	0.5	-169		Bi			54
092	V380 Ori NGC1999	05 34 00	-06 44 26	210 24	-19 43	0.48	-169	10	MR			140
						0.46		12.35				85
093	V380/OriNE	05 34 00	-06 41 00	210 21	-19 42	0.5 0.47	-168	12 20	Bi	0.04	7.0	3.2 ~ 3
094	V380/OriS	05 34 00	-06 56 00	210 35	-19 48	0.5	-169	13	MR			2-16
095	05341-0539 Ori A-E	05 34 11	-05 30 03	209 15	-19 07	0.5	-164	19	MR			40-400
096	AFGL5157 05345+3157	05 34 33	+31 57 40	176 31	+00 11	1.8 1.8	6	25 43.0	Bi	1.29	2.6	5.5E3 1.30E3
						1.8	19	8.0		0.58	1.61	1380
097	Ori-I-2 05355-0146	05 35 33	-01 46 50	205 57	-17 05	0.4	-118	8.0	Bi	0.16	2.3	12
098	05358+3543	05 35 49	+35 43 41	173 29	+02 26	1.8	76	25	Bi	1.8	1.6	6.3E3
						1.8	76	28				$10^{3.8}$
099	05358+3543 A+B ⁱ	05 35 13	+35 45 54	173 23	+02 21	1.8	73		Multi			~ 6300
100	05361+3539 G173.58	05 36 06	+35 39 06	173 35	+02 27	1.8	77	29.3	Bi	0.9	2.6	1.8E3
101	05363-0702 L1641C	05 36 19	-07 02 41	211 18	-19 30	0.5 0.48	-167	11.1	MB			17.6 40.2
102	Haro 4-255	05 36 57	-07 28 20	211 27	-19 24	0.50	-166	22,11	Bi	0.4	2.1	55
103	GGD4	05 37 21	+23 49 22	183 43	-03 40	1.0	-64	31	Bi	0.11	2.1	420
104	S235B	05 37 31	+35 39 55	173 43	+02 42	1.8	85	22	Bi			$< 1.3E4$
105	L1641S3 FIRSSE101	05 37 31	-07 31 59	211 34	-19 18	0.5	-165	13.7 36.8	Bi	> 0.37	> 2.0	61.5 1E2.9
106	L1641S	05 38 03	-07 28 59	211 35	-19 09	0.5	-164	28	Bi	0.75	3.8	294.7
107	L1641S4	05 38 25	-08 08 20	212 15	-19 22	0.5	-166		Bi	0.8		20.2
108	NGC2023-MM1	05 38 54	-02 19 36	206 52	-26 36			65	Bi			
109	NGC 2023 05391-0217	05 39 06	-02 17 24	206 51	-16 33	0.5	-142	13 27.9	Bi	0.27	1.2	889.1
110	NGC2024 FIR 4	05 39 12	-01 56 10	206 33	-16 21	0.415	-117		MR			
111	NGC2024 FIR 5	05 39 13	-01 57 03	206 33	-16 22	0.415	-117	30	MR			
112	NGC2024 FIR 6	05 39 14	-01 57 27	206 34	-16 22	0.415	-117	84	Bi			
	NGC2024 ^w							41,21				$\geq 10^{3.5}$
113	NGC2024 Ori B ^t	05 39 18	-01 56 42	206 34	-16 20	0.5 0.4	-141	60	Bi	0.51	3.7	4.6E4
114	L1641S2	05 40 23	-08 18 26	212 38	-19 00	0.5	-163	26.0	Bi	0.18	1.7	20.6 20.6
						0.46			Multi			200
115	05413-0104 HH212	05 41 19	-01 04 08	206 00	-15 29	0.46	-123	9.7	Bi	0.01	7.0	14
116	B35	05 41 42	+09 07 00	196 56	-10 25	0.46-0.5	-90	12.5	Bi	0.6	<u>2.3</u>	15
117	HH26IR	05 43 32	-00 15 23	205 04	-14 22	0.5	-124	30	Bi	0.46	<u>2.7</u>	50-300
118	HH25 MMS	05 43 34	-00 14 42	205 09	-14 36	0.5	-126	27	Bi			6
119	HH24	05 43 38	-00 13 23	205 31	-14 34	0.5	-126	15	Bi	0.36	<u>1.0</u>	20
120	NGC2068/LBS17	05 43 56	-00 01 42	205 23	-14 25	0.4	-94	15	Bi			< 2.0

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
121	NGC2068	05 43 58	−00 04 00	205 25	−14 25	0.5	−124	20	Bi			
122	CB34	05 44 03	+20 59 07	186 57	−03 50	0.6	−40	10	Bi	0.21	4.0	
123	NGC2071	05 44 30	+00 20 40	205 07	−14 07	0.5	−122	75,120	Bi	0.41	<u>3.7</u>	750
124	NGC2071N	05 45 00	+00 40 00	204 53	−13 51	0.5	−120	17	Bi	0.55	2.3	900 40-400 40
125	05487+0255	05 48 46	+02 55 02	203 19	−11 57	0.46	−95	12.3	Bi	1.30	3.0	7
126	05490+2658	05 49 05	+26 58 52	182 25	+00 15	2.1	9	20	Bi	0.71	1.5	4.2E3
127	05491+0247	05 49 09	+02 47 53	203 28	−11 55	0.46	−94	20	Bi	0.70	3.0	25
128	L1598NW	05 49 28	+08 20 48	198 35	−09 08	0.9	−143		Bi			
129	L1598	05 49 39	+08 12 55	198 43	−09 10	0.9	−143	14	Bi	0.22	2.8	
130	05553+1631	05 55 20	+16 31 46	192 10	−03 49	2.5	−166	17	Bi	1.89	1.4	6.5E3
	G192.16					2.0						3.26E3
131	CB39	05 59 06	+16 30 58	192 37	−03 02	0.6	−32	10	MR	0.5	2.0	
132	HD250550	05 59 07	+16 13 06	192 53	−03 11	1.0	−56	7	MR	1.5	<u>1.7</u>	150
133	S241	06 00 41	+30 14 54	180 52	+04 05	4.7	335	17.1	Bi	1.71	2.4	
134	06047-1117	06 04 21	−11 18 16	197 40	+04 29	0.5	39	24.5	Bi	0.26	3.3	6
135	Mon R2	06 05 22	−06 22 25	213 42	−12 36	0.8	−207	35	Bi	2.1	<u>3.0</u>	
						0.95		31				5E5
136	AFGL 6366S	06 05 41	+21 31 32	189 02	+00 47	1.5	20	18	Bi	0.99	1.7	2.0E4
137	AFGL 5180	06 05 54	+21 38 57	188 05	+00 53	1.5	23	19	MB	1.08	2.0	1.1E4
138	GGD 12-15	06 08 25	−06 10 45	213 53	−11 50	1.0	−205	25	Bi	0.87	<u>6.2</u>	1.0E4
139	S255-IRS1	06 09 59	+18 00 15	192 36	−00 03	2.5	−2	26	Bi	1.82	< 2	10 ⁵
	S255											
140	Mon R2E	06 10 22	−06 12 28	214 08	−11 25	0.95	−188		Bi			
141	06114+1745	06 11 29	+17 45 33	192 59	+00 09	1.75	5	20.0	Bi	1.28	1.2	4211
142	06291+0421	06 29 09	+04 21 44	206 50	−02 23	1.60	−66	20.0	Bi	0.81	1.2	1882
143	06308+0402	06 30 53	+04 02 27	207 20	−02 09	1.6	−66	20	Bi	1.44	1.4	3.9E3
144	AFGL961	06 31 59	+04 15 09	207 16	−01 48	1.6	−50	30	Bi	1.18	<u>1.2</u>	5.5E3
								40.3				
145	NGC2261	06 36 26	+08 46 57	203 46	+01 16	0.69	15	9	Bi	0.68	2.0	660
	RMON											
146	Mon OB1H	06 38 17	+10 39 56	201 18	+02 33	0.8	15	27.3	Bi	0.29		87
147	Mon OB1D	06 38 19	+09 37 32	203 14	+02 04	0.80	29	26,30	Bi	1.21	<u>1.9</u>	110
	NGC2264D											
148	Mon OB1I	06 38 19	+10 52 39	202 07	+02 39	0.80	37	18.2	Bi	0.55	<u>2.6</u>	
149	NGC2264G	06 38 26	+09 58 51	202 56	+02 16	0.7	28		Bi			~ 12
150	NGC2264	06 38 26	+09 32 00	203 19	+02 03	0.80	29	31.2	MR	0.24	<u>1.0</u>	2.3E4
	Mon OB1C											
151	Mon OB1G	06 38 27	+09 58 28	202 56	+02 16	0.80	32	54.6	Bi	0.89	<u>4.2</u>	< 450
152	S287N	06 45 19	−02 09 32	214 30	−01 49	1.4	−44		MB			
	06453-0209											
153	BFS 56	06 56 45	−03 50 41	217 18	−00 03	1.4	−1		Bi			
	06567-0350											
154	BiP 14	06 56 47	−03 55 28	217 23	−00 05	1.4	−2		Bi			
	06567-0355											
155	CMa-W	06 56 53	−11 54 46	224 29	−03 44	1.1	−72		Bi			
	06568-1154											
156	S287-B	06 57 06	−04 41 48	218 06	−00 22	1.4	−9		Bi			
157	S287-C	06 57 08	−04 36 10	274 37	−23 37	1.4	−561		Bi			
158	L1654	06 57 17	−07 42 16	220 47	−01 43	1.1	−33		Bi			
159	S287-A	06 57 55	−04 32 22	218 03	−00 07	1.4	−3		Bi			
160	Z CMa ^c	07 01 23	−11 28 36	224 36	−02 33	1.15	−51	14	Bi	0.15	1.4	

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
161	CB54	07 02 06	-16 18 47	228 60	-04 37	0.6	-48	10	Bi	0.78	5.0	
162	07028-1100	07 02 52	-11 00 05	224 31	-02 01	1.4	-49		Bi			279
163	L1660	07 18 01	-23 56 42	237 31	-04 50	1.0	-84	9	Bi			100
164	07427-2400 G240.31+0.07	07 42 45	-24 00 24	240 19	+00 04	6.6	8	40	Bi	1.6	1.5	4.9E4
165	WB89 1099 HH120	08 07 40	-35 56 07	253 18	-01 37	0.75	-20	17.7	Bi	0.32	1.7	46
166	WB89 1135	08 21 18	-41 46 10	259 37	-01 42	6.42	-190	21.5	Bi	3.3	2.4	1.4E4
167	HH46/47 ESO210-6A	08 24 16	-50 50 43	267 22	-07 30	0.45	-59	12.5 72	Bi	0.16	1.7	15.7
168	WB89 1173	08 39 23	-40 41 18	260 46	+00 41	1.35	16	19.3	Bi	0.61	1.4	480
169	WB89 1181	08 44 49	-43 43 28	263 47	-00 26	0.98	-7	25.9	Bi	0.33	1.2	1630
170	WB89 1187	08 47 01	-43 21 15	263 44	+00 07	2.24	4	25.3	Bi	1.86	1.43	7100
171	WB89 1189	08 47 39	-43 06 01	263 37	+00 22	0.59	4	15.4	Bi			310
172	08558-4347	08 58 49	-43 47 43	265 28	+01 28	0.7	18	30.0	Bi	1.08	1.1	
173	WB89 1275	09 22 46	-51 46 36	274 00	-01 09	6.38	-128	21.5	Bi	1.84	2.12	9.36E5
174	ϵ Ch I-C	11 05 28	-77 06 32	297 12	-15 41	0.14	-38	9.0	Bi			
175	ϵ Ch I-N	11 08 22	-76 18 06	297 01	-14 53	0.14	-36	12.3	Bi			> 19.5
176	BHR71 IRS2	11 59 00.	-64 52 02	297 43	-02 47	0.2	-9.7		MB ⁺			
177	11590-6452 BHR71 IRS1	11 59 03	-64 52 11	297 44	-02 47	0.2	-10	21	Bi	0.32	8.1	9
178	12091-6129	12 09 08	-61 29 37	298 16	+00 44	4.0	52	35.8	Bi	0.8	2.53	2.9E4
179	12326-6245	12 32 41	-62 45 57	310 08	-00 13	4.4	-17	35.5	Bi	0.6	1.8	3.8E5
180	12405-6238	12 40 35	-62 38 46	302 02	-00 04	4.5	-5	35.8	Bi			7.3E4
181	12496-7650	12 49 38	-76 50 45	303 02	-14 15	0.25	-62	18	Bi			80
182	12515-7641 HH52-53	12 51 00	-76 41 00	303 07	-14 05	0.25	-61	18	Bi			0.44
183	12522-7640 HH54	12 52 12	-76 40 01	303 11	-14 04	0.25	-61	14	MB			0.22
184	12553-7651 ^m	12 55 20	-76 51 22	303 22	-14 16	0.18	-44	3	Bi			1.2
185	13547-3944 CG12	13 54 42	-39 47 42	316 28	+21 05	0.63	144	12	Bi			110
186	14562-6248	14 56 13	-62 48 03	316 58	-03 42	0.7	-45	11	Bi	<u>0.4</u>		
187	Circinus 14563-6301	14 56 19	-63 01 42	316 52	-03 54	1.26 0.7	-86	30.0 19	Bi	4.5	2.0	73
188	14563-6250	14 56 18	-62 50 02	316 58	-03 44	0.7	-46	11	Bi	<u>1.0</u>		
189	14564-6258	14 56 25	-62 58 38	316 54	-03 52	0.7	-47	13	Bi	<u>1.4</u>		
190	14564-6254	14 56 29	-62 54 57	316 56	-03 49	0.7	-46	19	Bi	<u>1.0</u>		
191	14568-6304	14 56 52	-63 04 59	316 54	-03 59	0.7	-48	11	Bi	<u>0.25</u>		
192	14596-6320	14 59 37	-63 20 05	317 05	-03 39	0.7	-44	12	Bi	<u>0.2</u>		
193	14580-6303	14 58 05	-63 03 36	317 02	-04 01	0.7	-49	12	Bi	<u>0.4</u>		
194	14592-6311	14 59 17	-63 11 20	317 06	-04 12	0.7	-51	12	Bi	<u>0.8</u>		
195	16019-4903	16 01 59	-49 03 23	332 18	+02 17	2	80	40.0	Bi			4800
196	16191-1936	16 19 09	-19 36 25	356 05	+20 45	0.16	57	6.5	Bi	> 0.013	> 1.27	1.8
197	L1686 ^b	16 22 55	-24 12 06	353 04	+17 54	0.16	47	7.6	Bi			14
198	Gss30 EL21	16 23 22	-24 15 48	353 05	+16 17	0.16	47	9	Bi			11-12
199	ρ Oph AS	16 23 22	-24 18 07	353 03	+16 55	0.16	46	6.91	Bi			
200	ρ Oph A VLA1623	16 23 25	-24 17 46	353 04	+16 03	0.16	44	28	Bi	0.11	<u>14</u>	$\leq 5-10$

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
201	ρ Oph AN	16 23 28	-24 16 45	353 05	+16 55	0.16	46	16	Bi			
202	WL12	16 23 43	-24 28 04	352 59	+16 56	0.16	47	16.0	Bi			4.1
203	EL29	16 24 08	-24 30 37	353 00	+16 39	0.16	46	15.5	Bi			41
204	ρ Oph-B	16 24 13	-24 18 50	353 11	+16 45	0.16	46	12	Bi			0.5
205	IRS43	16 24 25	-24 34 10	353 01	+16 47	0.16	46	19.0	Bi	> 0.01	> 1.62	10.1
206	ρ Oph B2	16 24 26	-24 20 37	353 11	+16 42	0.16	46	13.28	Bi			
207	L1681B	16 24 26	-24 32 51	353 02	+16 33	0.16	46	13	Bi	0.02	3.3	13
208	IRS48	16 24 36	-24 23 55	353 10	+16 45	0.16	46	18.0	Bi			11.1
209	IRS51	16 24 38	-24 36 35	353 01	+16 33	0.16	45	11.0	Bi	0.05	2.89	1.3
210	L1709A ^d	16 27 50	-23 34 37	354 20	+16 34	0.15	43	4.2	MR			
211	L1709	16 28 34	-23 56 32	354 09	+16 38	0.16	45		Bi			
212	ρ Oph-S	16 28 54	-24 50 06	353 30	+15 29	0.16	43		Bi			
213	Re 13	16 28 52	-44 49 15	338 32	+02 37	0.90	33		Bi			288
214	V346N or FuOri	16 28 57	-44 49 08	338 33	+02 15	0.62	23	38	Bi	0.15	1.3	
215	ρ Oph-E L1689N	16 29 21	-24 22 13	353 56	+15 37	0.16	44	25	Multi	0.36	2.8	40-400 23
	16293-2422											27
216	16295-4452	16 29 33	-44 52 05	338 34	+02 01	0.6	21	13	Bi			35
217	16306-4455	16 30 40	-44 55 13	338 41	+01 50	0.7	22		MR	0.45	1.27	
218	16311-4457	16 31 07	-44 57 27	338 42	+01 45	0.7	21	11.0	Bi	> 0.48	> 1.43	
219	L43	16 31 38	-15 40 50	001 21	+20 58	0.16	57	9.5	Bi	0.32	<u>2.5</u>	4.3
	RNO91					0.16			Multi			4.3
220	16442-0930	16 44 14	-09 30 02	008 40	+22 11	0.16	60	3.8	MR			0.97
	L255											
221	L146	16 54 27	-16 04 48	004 30	+16 21	0.16	45		MR			
222	17081-2712 or B59-MMS1	17 08 10	-27 21 29	357 05	+07 08	0.16	20	17	Bi	0.58	2.9	
223	L100	17 13 04	-20 53 39	003 04	+09 58	0.23		9.5	Bi			
224	NGC6334	17 16 37	-35 54 49	351 10	+00 42	1.7	21	60	Bi			1.65E5
225	NGC6334B	17 17 18	-35 48 00	351 20	+00 39	1.7	19	24	Bi			4E5
						1.7						
226	NGC6334I ^c	17 17 34	-35 43 47	351 25	+00 39	1.74 1.74	20	175 80	Bi	0.31	2.0	8E4 8.0E4
227	CB81	17 19 19	-27 05 20	358 44	+05 16	0.6	55	6	MB			
228	17364-1946 L219	17 36 00	-19 46 00	007 00	+06 05			14	Bi			
229	W28-A2 G05.89-0.4	17 57 27	-24 03 54	005 53	-00 24	3.0 3	-21	140	Bi			4.2E5 3E5 3E5
230	H 36	18 00 36	-24 22 53	005 58	-01 10	1.5	-37	21	Bi	<u>0.07</u>		
231	M8E	18 01 49	-24 26 57	006 03	-01 27	1.5 3	-38	16	Iso	<u>0.31</u>	<u>1.0</u>	2.5E4
						5.7			Bi			
232	G9.62+0.19F ^m	18 03 16	-20 32 00	009 37	-00 12	5.7	-16		Bi			
233	18128-1943	18 12 50	-19 43 21	011 25	-01 22	1.8	-43	17.5	Bi			
234	18134-1942	18 13 24	-19 42 36	011 30	-01 29	1.8	-46	17.5	Bi			
235	18139-1952	18 13 59	-19 52 16	011 25	-01 41	1.8	-53		MB	1.77	2.0	
236	L483	18 14 51	-04 40 49	024 53	+05 23	0.2	23	15.5	Bi	0.1	3.6	14 9.5 ± 0.5
						0.2						
237	18150-2016	18 15 01	-20 08 10	011 11	-02 05	1.8	-66	9.5	Bi	1.94	2.8	
238	18151-1208	18 15 09	-12 08 34	018 20	+01 46	3.0	93	17	Bi	0.6	1.9	10 ^{4.3}
239	GGD27	18 16 13	-20 48 43	010 51	-02 35	1.7	-77	15	Bi	1.07	2.0	2E4
						1.7		18.5				2.0E4
240	18182-1433	18 18 17	-14 33 19	016 35	-00 03	11.7 4.5	-4	17	Bi	0.56	1.4	10 ^{5.1} 10 ^{4.3}

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
241	18196-1331 CRL2136	18 19 37	-13 31 47	017 38	+00 09	2.0	5	24	Bi			7.2E4
242	18258-0737	18 25 52	-07 37 30	023 34	+01 35	3.1	86	18	Bi			
243	18264-1152	18 26 27	-11 52 27	019 53	-00 32	12.5 3.5	-33	35	Bi	0.57	2.1	$10^{5.1}$ $10^{4.0}$
244	L379IRAS(3)	18 26 33	-15 17 51	016 52	-02 10	0.2 2	-8	7.0 48.8	Bi	0.44	1.6	> 40 $10^{3.8}$ $\sim 1.6E4$
245	L379IRS1 Serpens SMM1-11 SMM1-4,8,9?	18 27 18 18 27 25	+01 13 14 +01 12 16	031 35 031 35	+05 23 +05 20	0.31	29	28 65	Bi			84-210
246	S68 Firs1 SVS 4	18 27 25	+01 12 40	031 35	+05 21	0.5	47	28	Bi	1.11	<u>1.7</u>	300 90
247	L379IRAS(2)	18 27 43	-15 16 45	017 01	-02 24	2.0	-84	17	Bi	0.88	1.7	960
248	64 ^b , W4	18 28 40	-02 09 00	028 45	+03 31	0.7	37		MR	0.67	2.0	2.8E4
249	18316-0602 G25.65+1.05	18 31 40	-06 02 06	025 39	+01 03	2.8	51	67.4	Bi			1.76E4
250	18331-0035	18 33 08	-00 35 48	030 38	+03 15	0.31	18		Bi			2.5
251	18345-0641	18 34 35	-06 41 09	025 24	+00 06	9.5	17	24	Bi	1.4	1.8	$10^{4.6}$
252	18470-0044	18 47 02	-04 43 05	028 35	-01 44	8.2	-248	20	Bi	1.4	1.8	$10^{4.9}$
253	18537+0749	18 53 46	+07 49 30	040 30	+02 32	2.36	104	90 ^c	Bi	0.6	1.3	
254	G35.2-0.74	18 55 41	+01 36 30	035 12	-01 40	2	-26	28	Bi	1.39	2.6	1.6E4 1.2E4
255	18566+0408	18 56 41	+04 08 03	037 33	+00 12	6.7	24	34	Bi	1.0	2.4	$10^{4.8}$
256	R CrA	18 58 32	-37 01 30	359 56	-17 51	0.13	-40	28	Bi			96
257	19012+0536	19 01 18	+05 36 10	039 23	-00 08	8.6 4.6	-11	20	Bi	0.41	1.2	$10^{4.7}$ $10^{4.2}$
258	19035+0641	19 03 35	+06 41 55	040 37	-00 08	2.2	-5	22	Bi	0.37	2.4	$10^{3.9}$
259	W49	19 07 52	+09 01 11	043 10	+00 00	14.0	1	60	Bi			3E6
260	G45.07+0.13	19 11 00	+10 45 43	045 04	+00 08	6.4	15	24.4	Bi	1.2	2.2	$10^{5.77}$
261	G45.12+0.13	19 11 07	+10 48 11	045 07	+00 08	6.4	15	30	Bi	0.58	3.0	1.66E6
262	L723 2 Flows	19 15 42	+19 06 49	052 59	+03 03	0.3 0.3 0.3	16	20,60	Multi	0.4	<u>4.1</u>	2.8 2.4 6.5 3
263	CB188	19 17 57	+11 30 18	046 32	-01 01	0.6	-11	12.5	Bi	0.22	3.9	
264	L673 SMM1	19 18 04	+11 16 37	046 20	-01 09	0.3	-6	7.0	Bi	0.12	4.3	2.8
265	L673	19 18 08	+11 14 00	046 19	-01 11	0.3	-6	8	Bi	0.59	1.4	2.8
266	AS353	19 19 09	+10 56 30	046 10	-01 33	0.2	-5	17	Bi	0.27	<u>2.2</u>	7.2
267	19213+1723	19 21 23	+17 23 06	052 06	+01 03	4.12	75	28.0	Bi	1.10	1.44	2.82E4
268	19217+1651	19 21 44	+16 51 42	051 41	+00 43	10.5	132	37	Bi	1.6	3.4	$10^{4.9}$
269	L778	19 24 26	+23 52 37	058 09	+03 31	0.25	15	19	Bi	0.45	<u>2.3</u>	0.57
270	19266+1745	19 26 40	+17 45 41	053 02	+00 07	10.0	0.6	18	Bi			$10^{4.7}$ $10^{1.7}$
271	19282+1814	19 28 15	+18 14 32	053 38	+00 01	1.8	0.8	14	MB	<u>0.51</u>	1.5	9273
272	B335	19 34 34	+07 27 20	044 56	-06 33	0.40 0.25	-46	9	Bi	<u>0.4</u>	<u>2.8</u>	7.6 3
273	19374+2352	19 37 26	+23 52 57	059 36	+00 55	4.30	69	34.0	Bi	1.42	1.72	2.67E4
274	1548C27	19 40 48	+23 17 09	059 28	-00 03	2.4	-2	20	Bi			580
275	19410+2336(L)	19 41 04	+23 36 54	059 47	+00 04	6.08 2.48 6.4 2.1	7	65 ^c 44	Bi	2.4		$10^{4.0}$ $10^{5.0}$ $10^{4.0}$
276	19410+2336(U)	19 41 04	+23 36 54	059 47	+00 04	6.08 2.48 6.4 2.1	7	65 ^c 44	Bi			$10^{5.0}$ $10^{4.0}$ $10^{4.3}$ $10^{3.7}$
277	19411+2306	19 41 10	+23 06 47	059 22	-00 12	5.8 2.9	-10	28	Bi	0.52	3.0	< 970
278	L810 CB205	19 43 22	+27 43 40	063 36	+01 41	2.5	74	13	Bi	0.97	1.4	
279	S87	19 44 14	+24 28 00	060 53	-00 08	2.7	-6	20	Bi			3.5E4
280	S88 S88 B	19 44 40	+25 05 30	061 28	+00 06	2.0 2.0	4	17 23.5	Bi	0.9	1.5	1.8E5

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
281	19471+2641	19 47 06	+26 41 16	063 07	+00 27	5.06	39	20	Bi	6.8	> 1.7	
282	19520+2759	19 52 04	+27 59 30	064 49	+00 10	2.62	10	30	Bi			1E5
283	19529+2704	19 52 56	+27 04 54	064 08	-00 28	3.37	-28	21.5	Bi	1.03	1.9	3.9E4
284	19550+3248	19 54 59	+32 48 26	069 15	+02 09	4.04	75	14.9	Bi			600
285	K3-50	19 59 50	+33 24 19	070 18	+01 36	2.0		15.5	Bi	2.6	2.5	
	19598+3324					8.8	246	23				
286	CB214	20 01 54	+26 29 42	064 42	-02 29	0.6	-26	5	MB			
287	CB216	20 03 45	+23 18 25	062 13	-04 32	0.6	-48	16	Bi			
288	20050+2720	20 05 02	+27 20 09	065 47	-02 37	0.7	-32	140	Multi			260
289	20056+3350	20 05 36	+33 50 53	071 19	+00 50	1.67	24	58.0	Bi	0.43	2.24	4E3
290	CB217	20 05 55	+36 58 14	073 58	+02 28	0.6	25	5	MR			
291	20062+3350	20 06 17	+33 50 32	071 23	+00 42	0.08	2	1	Bi	0.02	1.48	5.4
292	20106+3545	20 10 38	+35 45 42	073 29	+01 01	1.64	29	15.0	Bi	0.46	1.18	1.79E3
293	20126+4104	20 12 41	+41 04 20	078 07	+03 38	1.7	108	63.1	Bi	1.35	<u>1.7</u>	(1-6.3)E4
								48.9				
							4.18	61				7.12E4
294	20188+3928	20 18 51	+39 28 18	077 28	+01 46	2.0	61	18	Bi	1.4	1.6	$10^{4.10}$
						0.3		36				1.26E4
295	G75N	20 19 50	+37 17 42	075 47	+00 21	5.5	34	40.3	Bi	0.58	3.3	330
296	G75C	20 19 52	+37 17 04	075 47	+00 20	5.5	33	33.6	Bi	0.56	1.6	5E3
297	G75E	20 19 55	+37 16 52	075 47	+00 20	5.5	32	42.9	Bi	0.60	3.4	430
298	G75.78NE	20 19 53	+37 27 22	075 56	+00 26	5.5	42	36.4	Bi	1.1	1.5	$\sim 10^4$
299	20216+4107	20 21 38	+41 07 56	079 08	+02 17	1.7	68	15	Bi			$10^{3.3}$
300	V1515Cyg	20 22 03	+42 02 40	079 55	+02 44			11.5	MB			
301	20227+4154 ^g	20 22 46	+41 54 29	079 53	+02 33	2	89	5.3	Bi	1.7	1.8	$10^{3.46}$
302	20228+4215 ^g	20 22 50	+42 15 13	080 10	+02 45	2	96	16.9	Bi	1.3	1.4	$10^{3.41}$
303	20231+3440 SMM1	20 23 04	+34 40 16	074 01	-01 42	1.0	-30	31	Bi	1.2	2.6	
304	20231+3440 SMM2	20 23 05	+34 43 16	074 04	-01 40	1.0	-29	31	Bi			
305	20272+4021 ^g	20 27 17	+40 21 54	079 08	+00 58	2	34	7.3	Bi			$10^{3.4}$
306	AFGL2591	20 27 36	+40 01 14	078 53	+00 43	2.0	25	74	Bi	0.68	1.9	9E4
	20275+4001					0.7						$10^{3.96}$
307	20281+4006 ^g	20 28 12	+40 06 29	079 02	+00 40	0.7	23	7.3	Bi	1.50	1.5	$10^{2.5}$
308	20286+4105	20 28 41	+41 05 39	079 53	+01 11	3.7	14	14.0	Bi	1.35	2.54	3.9E4
309	20293+3952#1	20 29 21	+39 52 59	078 58	+00 21	2.0	8	41	Bi	0.22	3.5	$10^{3.8}$
						1.3						$10^{3.4}$
310	20293+3952#2	20 29 21	+39 52 59	078 58	+00 21	2.0	8	41	Bi			$10^{3.8}$
						1.3						$10^{3.4}$
311	20343+4129#1	20 34 20	+41 29 33	080 50	+00 34	1.4	14	7	Bi	1.34	1.6	$10^{3.5}$
312	20343+4129#2	20 34 20	+41 29 33	080 50	+00 34	1.4	14	7	Bi			$10^{3.5}$
313	20353+6742	20 35 19	+67 42 30	102 20	+15 58	0.44	121	9.5	Bi			3.3
	L1152									0.36	3.5	
314	W75-N	20 36 51	+42 27 20	081 53	+00 47	2.0	27	22	Bi	0.78	<u>3.5</u>	
						2.0						
								62.5				1.4E5
315	DR21	20 37 13	+42 08 50	081 41	+00 32	2.0	19	27	MB			6E5
						3.0	60	53				1.5E5
316	L1036	20 37 43	+56 58 56	093 34	+09 30	0.44	72		Bi			
317	L1157	20 38 40	+67 51 33	102 39	+15 48	0.44	120	15.5	Bi			11
318	PVcep	20 45 23	+67 46 36	102 58	+15 14	0.5	131	10	Bi	0.22	1.6	80
319	20520+6003	20 52 05	+60 03 18	097 11	+09 54	0.44	76	12.0	Bi	0.25	~ 3.4	3.0
	L1082A											
320	20526+5958	20 52 41	+59 58 19	097 10	+09 48	0.44	75	8.5	Bi	0.40	~ 3.3	0.8
	L1082B											

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
321	V1057Cyg	20 57 06	+44 03 47	085 28	-01 03	0.7	-13	15	MB	1.14	3.1	540
322	20582+7724	20 58 12	+77 24 05	111 40	+20 13	0.1	34	15	Bi	0.68	<u>1.8</u>	0.4
323	V1331Cyg	20 59 32	+50 09 56	090 19	+02 41	0.7	33	20	MR	0.83	1.0	36
324	L988-a	21 00 45	+49 51 13	090 13	+02 19	0.7	28		Bi	2.05	2.6	19
325	21015+6757	21 01 32	+67 57 15	104 05	+14 09	0.44		11.5	Bi			134.6
326	L1172D	21 01 45	+67 42 32	103 54	+13 58	0.44	106	15	Bi		<u>1.8</u>	0.95
327	L988-e	21 02 20	+50 02 40	090 31	+02 16	0.7	282		Bi			140
328	L988-f	21 02 25	+49 55 50	090 27	+02 11	0.7	26		Bi	2.23	> 1.92	28
329	B361 ^b	21 10 41	+47 12 01	089 23	-00 41	0.35	-4	9.1	Bi			4.7
330	L944 SMM1	21 15 51	+43 06 08	087 04	-04 11	0.7	-51	11.0	Bi	0.18	3.1	4.4
331	CB230	21 16 55	+68 04 52	105 10	+13 09	0.6	136	8.2	MR			
332	IC1396-W	21 24 39	+57 43 14	098 20	+05 14	0.75	68		Bi	1.87	1.7	
333	21307+5049	21 30 47	+50 49 03	094 16	-00 25	4.94	-359	20.0	Bi	0.09	2.2	1.23E4
334	21334+5039	21 33 24	+50 39 43	094 28	-00 48	5.0	-70	23	Bi			2.5E4
335	CB232	21 35 14	+43 07 05	089 38	-06 37	0.6	-69	11	Bi	0.22	2.8	
336	V645Cyg	21 38 11	+50 00 43	094 36	-01 48	5.6	-176	26	Bi	1.87	1.7	1.1E5
						6.0		28				
337	GN21.38.9	21 38 54	+56 21 33	098 51	+02 55	0.75	38	10	Bi	0.09	2.2	(1-1.1)E2
338	IC1396-N BIMA1	21 39 08	+58 02 24	099 59	+04 10	0.75	54	19	Bi			
339	IC1396-N ^u	21 39 10	+58 02 29	099 59	+04 10	0.75	54	20	Bi	0.38	2.9	(5-3.4)E2
								42.9				2.6E2
340	IC1396-N BIMA2	21 39 10	+58 02 31	099 59	+04 10	0.75	54	43	Bi			
341	1413+5442 or G98.04	21 41 21	+54 42 32	098 02	+01 27	7.4	187	16.9	Bi	1.1	1.3	(2.5-13.9)E4
342	NGC7129 FIR ^u	21 41 52	+65 49 50	105 22	+09 51	1.0	171	13	Bi	1.1	<u>1.9</u>	(2-10)E2
343	NGC7129 FIRS2-MM1	21 41 52.5	+65 49 36	105 22	+09 50	1.25	171		Bi			10
344	NGC7129 FIRS2-IR	21 41 52.6	+65 49 40	105 22	+09 50	1.25	171		Bi			
345	NGC7129 FIRS1-IRS6	21 41 57.3	+65 53 08	105 25	+09 53	1.25	172		Bi			500
346	NGC7129 FIRS1-MM1	21 41 57.3	+65 53 10	105 25	+09 53	1.25	172		Bi			50
347	LKH α 234	21 41 57	+65 53 05	105 25	+09 53	1.0	172	15	MR	0.8	<u>1.3</u>	1E3
348	21428+4732	21 42 51.2	+47 32 55	093 35	-04 10	1	73	11.0	Bi	0.66	2.5	< 4
349	21429+4726	21 42 59	+47 26 40	093 32	-04 16	1.0	-74	13	Bi	0.6		35
								14.5				35
350	21432+4719	21 43 17	+47 19 30	093 29	-04 23	1.0	-76	15	MB	1.4-2.1		40
								15.0				40
351	21441+4722	21 44 06	+47 22 11	093 38	-04 26	1.0	-77	13.5	Bi	1.2		< 5
								14.0				< 5
352	IC1396-E	21 44 31	+57 12 29	099 59	+03 04	0.75	40		MR			96
353	EL1-12	21 45 27	+47 18 08	093 46	-04 38	0.9	-73	24.7	MR	0.23	<u>1.1</u>	95
	L1031B							25				245
	IC5416							29				
								26.1				157
354	21461+4722	21 46 07	+47 22 40	093 54	-04 39	1.0	-81	9	MR			11
								9				11
355	BBD+46°3471	21 50 39	+46 59 34	094 15	-05 26	0.9	-85	15	Bi			150
356	21519+5613	21 51 58	+56 13 34	100 10	+01 40	7.30	212	30.0	Bi	2.49	1.89	1.91E4
357	22051+5848	22 05 10	+58 48 06	103 08	+02 39			8.7	Bi			
	L1165											
358	22103+5828	22 10 24	+58 28 42	103 30	+02 00	0.90	31	16	Bi	0.25	1.2	67
359	22134+5834	22 13 24	+58 34 12	103 53	+01 51	2.6	29	26	Bi	1.25	1.48	10 ^{4.1}
						0.9		19.0				1580
360	22142+5206	22 14 15	+52 06 26	100 23	-03 35	4.5	-281	20.2	Bi	2.1	3.6	16100

Table 1a CO Molecular Outflows(continued)

No.	Name	α h m s	δ ° ' "	L ° ' "	b ° ' "	D kpc	Z pc	ΔV km/s	Po.	$R_{\max}$ pc	R_{coll}	L_{bol} $L_{\odot}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
361	22172+5549	22 17 18	+55 49 40	102 48	-00 43	2.87	-36	18.0	Bi	1.48	1.64	4.78E3
362	S140	22 17 41	+63 03 49	106 48	+05 19	1.0	93	42	Bi	0.24	1.7	2E4
						0.91		120				1.4E4
						0.9						5E3
						0.9						(0.5-9.0)E4
363	S140-N	22 17 52	+63 17 50	106 57	+05 30	0.9	86	32	Bi	0.21	1.4	40-400
	22175+6317											
364	L1204-A	22 19 51	+63 36 33	107 18	+05 38	1.28	88	34	Bi			1.24E3
	22198+6336					0.9						
365	L1204-B	22 19 56	+63 22 12	107 11	+05 26	0.9	85		MB			
	22199+6322											
366	L1221	22 26 37	+68 45 52	110 39	+09 38	0.2	33	14.1	Bi	0.31	2.1	2.7
	22266+6845											
						0.2			Multi			2.7
367	L1203	22 26 47	+62 44 22	107 30	+04 29	0.9	70		Bi			
	22267+6244											
368	L1206	22 27 12	+63 58 21	108 11	+05 31	0.9	86	15	MB	0.13	2.1	1.2E3
369	22305+5803	22 30 31	+58 03 30	105 30	+00 14	5.40	22	26.0	Bi	2.14	2.09	1.41E4
370	22336+6855	22 33 36	+68 55 20	111 17	+09 27	0.58	49	12	Bi			0.6
371	22343+7501	22 34 22	+75 01 32	114 29	+14 42	0.20	51	7	Bi	0.55	1.6	27
372	22376+7455	22 37 41	+74 55 50	114 38	+14 30	0.2	50	< 6	Bi	<u>0.24</u>		14
	L1251											
373	L1211	22 45 23	+61 46 07	108 55	+02 35	0.75	34		Bi			
	22453+6146											
374	22475+5939	22 47 31	+59 39 03	108 12	+00 35	4.7	48		Bi	1.6	1.8	5.1E4
	S146					5.2		18.0				6.2E4
375	22506+5944	22 50 37	+59 44 50	108 35	+00 30	5.1	44	10.5	Bi	0.94	1.2	1.5E4
	10860+0049											
376	Cep A	22 54 19	+61 45 44	109 52	+02 07	0.73	27	40	Bi	0.5	<u>6.2</u>	5E4
								132				2.5E4
												1.9E4
377	22570+5912	22 57 03	+59 12 22	109 06	-00 21	5.1	-31	24	Bi			10 ^{4.7}
378	23011+6126	23 01 10	+61 26 16	110 29	+01 29	0.73	20	20	Bi	0.5	3.8	50
	Cep E											
379	23032+5937	23 03 17	+59 37 40	109 60	-00 17	3.5	-17	20	Bi			1.1E3
380	23033+5951	23 03 19	+59 51 59	110 06	-00 04	3.5	-4	34	Bi	0.90	2.6	10 ^{4.0}
381	Cep C	23 03 46	+62 13 49	111 04	+02 05	0.75	27		Bi			
	23037+6213											
382	MBM 55	23 05 54	+14 49 00	089 11	-40 56	0.18	-118	6.5	Bi	0.44	2.2	
383	NGC7538 ^f	23 11 36	+61 11 49	111 32	+00 47	2.8	38	35	Iso	1.3	1.0	1.6E5
384	NGC7538	23 11 36	+61 10 33	111 32	+00 45	2.7	36	30	Bi	0.4	1.0	1E4
	IRS11					2.8						(0.69-1.1)E4
385	NGC7538	23 11 37	+61 11 48	111 33	+00 47	2.7	36	18	Bi	0.5	1.1	1.5E5
	IRS1					2.8						(5.1-13.6)E4
386	NGC7538 A ^d	23 11 39	+61 10 18	111 32	+00 45	2.7	35	30	Bi	0.65	3.3	
387	NGC7538	23 11 45	+61 10 48	111 33	+00 45	2.7	36	30	Bi	0.8	1.0	3E4
	IRS9					2.7						(4.7-5.2)E4
388	23139+5939	23 13 58	+59 39 00	111 15	-00 46	3.5	-47	18.5	Bi	0.7	1.3	1.9E4
	G111.25					3.5						2.5E4
						4.8		28				2.5E4
389	23140+6121	23 14 02	+61 21 22	111 52	+00 49	6.4	92	17	Multi			4.35E4
390	23151+5912	23 15 09	+59 12 25	111 14	-01 14	5.7	-76	54	Bi	0.47	1.3	1.0E5
						3.5						3.9E4
391	MWC1080	23 15 16	+60 33 52	111 44	+00 02	2.5	1	26	Iso	1.2	<u>1.0</u>	1.7E4
								109				
392	L1246 SMM1	23 22 52	+63 20 10	113 31	+02 20	0.73	30	6.0	Bi	0.04	2.2	1.8
393	CB244	23 23 48	+74 01 08	117 07	+12 24	0.2	43	9.1	Bi	0.03	1.5	0.9
	L1262					0.6				0.39	3.7	
394	23314+6033	23 31 25	+60 33 55	113 36	-00 37	2.78	-30	46.0	Bi	0.22	1.74	1.09E3
395	23385+6053	23 38 28	+60 53 51	114 32	-00 33	4.7	-46	8.6	Bi	0.85	1.3	24300
						6.9		24				7.57E4
396	23545+6508	23 54 34	+65 08 29	117 19	+03 09	1.4	77	26.2	Bi	0.9		4E3
397	CB247 ^b	23 55 04	+64 30 10	117 14	+02 30	0.6	26	11	MR			

Table 1b Parameters of CO molecular outflows

No.	Name	M $M_{\odot}$ (16)	P $M_{\odot} \text{ km/s}$ (17)	E 10^{45} erg (18)	t 10^4 yr (19)	L_m $L_{\odot}$ (20)	F $10^{-4} M_{\odot} \text{ km/s yr}$ (21)	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$ (22)	IIH2WI ^a (23)	Ref. (24)
001	LKH $_{\alpha}$	1.5 ^h	2.6	0.056		0.02	0.4		H W	1,7,347
002	00117+6412	45.5	321.8	8.00E+02	8	8.2	0.07 40	0.32 4	W	2,251 ^k 382,345 207
003	00213+6530	1.13 ^h	8.31	0.611	18	2.94E-2			W	85,347
004	00259+6510	1.52 ^h	14.2	1.32	11.5	10.01E-2				85
005	00259+5625 CB3	0.4 4	2.8 37.2	0.2 5.5	1.5 0.5-1	0.11 5.6	2 53	0.2 32	WI	240,231 201,178,179
006	00338+6312 L1287 RNO1b/1c	0.3 2.5 4.5 ^h	4.5 13.2 51.4	0.69 0.79 7.3	1.2 11	0.5 0.06		0.15	W	104,345 3 118
007	00342+6374	11	385	133	2.8	39	140	0.4		370
008	00420+5530	46.9	238.7	17.7					W	207,346
009	00494+5617	11.4	59.2	3.4	24	0.12	2.5		WI	3,349,345,424
010	NGC281-E									86
011	01133+6434	2-5	7-19	0.6-2	8	0.06-0.2	0.9-2		W	87,345
012	W3-IRS5				> 0.31				II W	1,4,341,345
		8.02 2016	110.1	16.07	3.6	2594	7560	4.8		152,255,348 239,349
013	IC1805-W								H W	86,254,328,347
014	02310+6133	508.1	2.52E+03	1.00E+02	51.5	1.59	31	2.45		381
015	02461+6147	85	271	30	35	0.71	8	0.26	II	380
016	AFGL4029 02575+6017	9.7	33.1	1.2	17	0.06	1.9		IIH W	32,337,346
017	AFGL437	> 23 ^h 620		20	6 5.77	2.7 750	28 2350		II 2W	1,5,7,345 239,302,337
018	RNO13	0.2			2.3			3.4E-2	H2	7,8,265,266
019	L1448 IRS2								IIH W	242,345,347,374
020	L1448 IRS3	0.07 0.06	1.9	0.66	0.32 0.56	0.52	1.7	2	H2WI	9,232,265,266,410 228,345,347,374
021	L1448 U-star	0.06				2			H2WI	9,266,232,347,410
022	AFGL490-iki 03234+5843	3.4		27	2				II	192,274
023	AFGL490	15.4 ^h 4.9-14 30	1.33E+02 44.3	14 68-150 200	1.7 1.8 1.2	6.8 115,19	78 400,76 600	1.3 ^c	II 2W	11,252,268,302 1,7,152,228 10,345
		15.4	493	158	1.8	19				255,354
024	L1455NW	0.3-0.9 0.3-0.9				0.01-0.1 0.01-0.1	0.3-2.8		H W	13,345 14 12
025	RNO15FIR L1445 FIR	0.07 0.2			2.2			9.2E-3	H2W	8,268 14,345 109
026	L1455M	0.6-3.1	2.9-30.2	0.2-3.1	4.7-2.9	0.04-1.0	0.6-9.8	0.06-0.98	H W	14,8,13,108,228,345
027	NGC1333								H W	16,347
028	IRAS2NNE-SSW	0.251	1.38	0.106	1.7	0.011	0.19		H2W	222,267,345,347
029	IRAS2east-west	0.0287	0.165	1.29E-2	0.9	0.0157	0.24		H2W	222,267,345,347
030	HH12	0.1 0.2 0.09	0.6 1.2 0.31	0.04 0.08 0.016	0.7 1 2.7	10 ^{-1.2}	10 ^{0.1}	3.0E-2 0.04	H2WI	17,297,334,347 18,345 222,90
031	HH7-11 SSV	4 1.03 4.2 ^h 2			6.4 2.6 1 2 1		40 12 17		H2W	7,135,271 228,345 19,347 96,392
		0.3 0.034	11 0.201	4 0.0152	3.3 1.2	0.47 0.0123	4.5 0.19	20		222 222,169,135,267 232,231,331,345,390
032	NGC1333 IRAS4A								H2WI	
033	HH6	0.038	0.173	0.0105	0.57	0.0162	0.31		H W	222,345
034	03271+3013				~ 0.1				H2	121,269
035	03282+3035	0.09 0.06	~ 2.2	~ 0.77	0.75 1.3 ^s	0.9 0.13	3 0.56	1.19	2	121,269,270 228
036	03301+3057 Barnard 1	7E-4 5E-3	0.01	1.8E-3	≤ 0.1	1.6E-2			H W	126,418 200
037	HH211	0.02			< 0.1				H2	209,278,407
038	B5-IRS3	0.1	0.4	0.02	9				H2	21,20,272
039	B5-IRS1	0.02	0.3	0.05	1				H2	21,20,272,391
		1.2-50 5	12.6-230 16	1.4-11.7 0.39	10 24	0.26		10		272,224,310 21,20

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot}\text{km/s}$	E 10^{45}ergs	t 10^4yr	L_m $L_{\odot}$	F $10^{-4}M_{\odot}\text{km/s yr}$	$\dot{M}^o$ $10^{-5}M_{\odot}/\text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
041	B5-IRS2	0.4	2	0.15	14					21,20
042	L1489	0.04	0.04	1.2E-3	13	4E-4	0.002	2E-4	IIH2WI	22,13,23,274,347,375
043	PP 13S	~ 0.01	0.24		0.61		~ 0.15		H W	323,345
044	04166+2706						0.031		H I	190,330,334
045	04181+2655						0.022		H	190
046	T Tau	> 0.10				> 0.068	> 0.6		H W	7
		0.22			4			1.4E-2		80,12
		0.44			0.98	0.17	1.7			228
047	04191+1523				~20		< 0.02		I	86
	IRAM04191	0.03								146,367,368
048	04239+2436	4.3	> 3.2	> 0.026	0.4	(0.11-3.6)E-2			2	383
049	Haro 6-10								H I	238,145,146,274
	L1524									84,86
050	ZZ Tau	0.04	0.11	0.3E-2	4	7E-4	3.1E-2	3.1E-3	H	24,13
051	L1551-IRS5	0.3		6.7E-2	3	0.2	2	8E-2	H2 I	1,25,26,267,334
		1.22	5.6	0.36	3	0.3	5			245,276
		3.11			4.7	2.1	11			228,402
052	L1551NE	0.34	0.68	2.75E-2	~ 60				H I	128,232
		6.9E-3	0.03	1.5E-3	~ 0.2	0.05				184
053	HL/XZ Tau	0.01			0.7			4.1E-3	H W	8,243
		0.001	0.003	0.0001	1.6		2.1E-3	6.0E-3		80,347
054	TMC2A ^b									23
										172
055	L1529	< 1							H	27
056	04325+2402	0.06	0.17	0.5E-2	19	3E-4	9E-3	0.9 E-3	2W	24,22,31,215
	L1535									13,347
057	L1642	0.0004-0.01 ^h	0.001-0.02	0.2-6E-4	3.6		0.0002-0.005	0.04-1E-3	H	29
058	L1527	0.1-0.5						2E-3	I	30,231,232,235,253
	L1527 IRS	0.176 ^c			1.2	0.163	1.61	1.77		246,367,331,329
059	04361+2547	0.008 ^h			0.6		0.04		WI	144,345
060	TMC1A	1.5E-2 ^h	1.1E-1	9.2E-3	0.227	7.2E-2	0.64		I	197,31,84,145,330,334
061	IC2087	0.28	0.72	2.0E-2	19	8E-4	0.038	3.8E-3	H	24,13
062	04381+2540								2	84,145,274
	TMC1	9.5E-3 ^h	7.1E-2	6.7E-3		0.038	0.29			197,146,23
063	05137+3919	115.7	724.7	76.7					W	207,346
064	HH114/115	0.3								383
065	05168+3634	15.6	247.8	50					W	207,346
066	L1634								H2	86,25,215,267,268
067	AFGL5142	4.9	14.3	0.6	20	0.03	0.72		H W	32
	05274+3345	1082			3.28	5.47	109			239
		3.1	170		1.7	5.8		18		313
		2.4	63		1.3	2.7		18		313
068	L1582B	0.12-0.6	0.42-2.1	0.016-0.081	9.5	1.4-7E-3	0.044-0.22	0.0044-0.022	H I	31,28,232
069	RNO43 ^l	24			20	0.67	12		H I	228,232
		0.42 ^{hv}	1.63	0.74	0.13	6	4	0.4		351
070	RNO43S	2.1	5.8	0.4	10			3E-2	H	17,7
071	RNO43N	2.2	5.6	0.4	8			4E-2	H	17,7
072	Ori A-W	16 ^h		1.9		0.037			W	86,346
										33
073	HH83	0.11 ^h	0.438	0.0187	5	3.2E-3	0.092		H	165
074	Orion-S	2.8			4			0.21	H W	8,12,103,35,37,34,349,345
										349,345
075	CSO 2	0.09	0.22		2.3		9.60E-02	0.39	H	262
076	05327+3404								H	371
										395
077	Orion-A	8.2 ^h	160	42	0.15	230	1100	100	IIH2W	11,151,37,41
	Orion-KL	5		2.00E+02		2600	5200			1,7,39,249
		8.2	533	346	0.075	1.00E+03				255,345,349,278
		8.2			0.075	1.00E+03	2.00E+03			228
078	Orion-KLN	0.0016		> 1.2E-2					H W	129,349
079	AC 3	0.24	1		3.3		0.31	0.73	H	262,404
080	FIR 1	0.49	1.4		1.7		0.8	2.9	H W	262,345

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot}\text{km/s}$	E 10^{45}ergs	t 10^4yr	L_m $L_{\odot}$	F $10^{-4}M_{\odot}\text{km/s yr}$	$\dot{M}^o$ $10^{-5}M_{\odot}/\text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
081	OMC-2/3(MMS 9)	0.6	3.2		3.3		0.98		H2W	262,345,345
		12.3	30.9	1.22	5		266			316,404
082	OMC-2	1.8 ^h	6.5	0.33	0.9				H2W	88,316,277,349,345,404
083	HH 34	0.0009	0.004	1.63E-04						261
084	MMS 10	0.29	1.3		3.1		0.42		H	262
085	L1641N	1.6 ^h	9.3	0.57	1.4	0.15			H2W	133, 33,34,273
		2.18 ^h	49	11	0.99	2.0-9.1	10-49			116,347,291
086	CS-star HH1	1.1E-3	0.0458	1.994E-2	0.1				H2W	195,7,137,275,278
	VLA1	0.7			1			0.22		8,345,386
										34
087	05339-0647								H W	263,345
088	VLA3	0.0987	1.2029	0.177					H W	195,263,345
089	VLA 2								H W	263,345
090	HH 1-2	0.073	0.35	1.56E-2						261
091	05339-0646	3.5	32	2.4	4	0.6	9	0.9		421
092	V380 Ori	0.8			3			0.064	H2W	8,34
	NGC1999	1.9	8.4		9			0.05		17,278,345
093	V380/OriNE	0.32			4			0.024	H2	8,7,273
		0.2	2.6	0.36	1.58	0.2-1	1.6			264
094	V380/OriS	2.8			4			0.21	H	8,7,12
095	05341-0539 Ori A-E	21 ^h		4.7		0.24			W	86,33,346
096	AFGL5157	7.6	37.1	2	19	0.09			H2W	32,282,3
	05345+3157	672			2.49	217	1310			239,203,346
		6.1	38.3	3.7						207
097	Ori-I-2	0.5 ^h	0.2	0.05	5	6E-3			H	89
	05355-0146									
098	05358+3543	15.7	87.8	5.4	17	0.26	5.2		2W	3,279,314,350
		8.9 ^c	65	4.7	3.25	1.2	7.8			203,345
		21 ^h	288	40	3.7	9.1	79			237
099	05358+3543 A+B [†]	9.6			3.7				W	325,345
100	05361+3539	32	170	9.4	5.3	1.4			II 2W	194,255,284,345
	G173.58									
101	05363-0702	2.6		0.19		0.006			2	34,86,278
	L1641C				11	0.006				138
102	Haro 4-255	0.72			5			0.042	H2W	8,34,153,278,291
										326 [†] ,161,347
103	GGD4								W	90,345
104	S235B	> 2 ^h							II W	1,6,7,77,341,349,345
105	L1641S3	0.35		0.17		0.017			2W	86,269,278
	FIRSSE101	0.38	7	1.3	0.99	<3.2-1.1	<2.1-7.0			116,345
										34
106	L1641S	1.4		0.3	7	0.013			H2	138,33,86,269
107	L1641S4	0.08		0.23	10	0.003				86,138
108	NGC2023-MM1	0.52 ^c	3.5	0.48	0.25	0.29	30.3			218
109	NGC 2023									59
	05391-0217	2.86 ^h	22.26	1.6	2.9	0.44	7.6	0.76	IIH2W	181,1,7,182,281,374
110	NGC2024 FIR 4	0.04	0.4	0.04	0.26	0.13				360,388
111	NGC2024 FIR 5	1.6 ^h	20	3.4	1.4	2			WI	107,360,364,232,388,418
112	NGC2024 FIR 6	0.13	2.1	0.7	0.04-0.33	15			I	360,232
		0.13	2.1	0.7		15				102
	NGC2024 ^w	0.59	8.53	1.35						388
113	NGC2024 Ori B [†]	5.5 ^h	82.5	25.3					II 2WI	101,1,7,283,337,338
										349,345,346
114	L1641S2				2.5	0.037				138
		0.8		0.37		0.037				86
		0.65 ^h	4.1	0.35	0.7					99
										383
115	05413-0104	0.2							H2W	215,278
	HH212									345
116	B35	0.86	2	6.4E-2	18	2.9E-3	0.11	0.011	H	22,28,13,315
117	HH26IR	3.1				1.8	15		H2WI	7,280,232,331
		3.1	13	2.4	3		4.3	0.22		43
118	HH25 MMS	0.05 ^c	0.16	6.7E-3	1.7	7.1E-3	9.2E-2		IIH2 I	408,232,409
119	HH24	3.2				0.24	4		H2WI	7,280,345
		3.2	15	1.7	6		2.7	0.13		43,387,409
120	NGC2068/LBS17	0.12	0.71	0.052	1.8		0.394		H	379

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot} \text{ km/s}$	E 10^{45} ergs	t 10^4 yr	L_m $L_{\odot}$	F $10^{-4} M_{\odot} \text{ km/s yr}$	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
121	NGC2068	2.1	6.9	0.5	20			8E-3	H	17,67
122	CB34	0.6	4.2	0.29	1.5	0.15	2.8	0.28	H2	240,394
										178,179
123	NGC2071	20		2.80E+02		175	600		IIH2W	1,7,45,175,266,376
		9.7	81.4	8	1.6	4	51	4		11,46,152,297,345,374
		9.7			1.4	24	91			228,349,346
124	NGC2071N								H	33
		5		0.3-3		0.05-0.2				110
125	05487+0255	0.8 ^h	21		3.7	1.2			H	75
126	05490+2658	7.1	25.5	0.97	37	0.02	0.69			3
127	05491+0247	0.2 ^h	2.7		5.4	0.055			H2 I	215,75,114,333,411,412
128	L1598NW								H	86
129	L1598	0.5			2.2	49	1.6		H	91
130	05553+1631	11.9	44.3	2	46	0.04	0.96	0.035-0.26	IIH2W	3,307,255,339,413
	G192.16	58	210	9.4	27	0.2	7			194,255,346
131	CB39				30					178,179
		0.6	4.8	0.4	2.1	0.14	2.2			240
132	HD250550	> 1.8				> 0.03	0.6			7
133	S241	64.99	554	40	14	2.4-16	40	4	W	198,1,346,345
134	06047-1117								H2	220
135	Mon R2	100 ^h		2.30E+02		24	200		II W	1,7,255,338
		202	~ 2000	290	15	16	160			81,255,349,345
136	AFGL 6366S	3.4	20.1	1.3	11	0.1	1.8		II W	32,337,346
137	AFGL 5180	2.3	12.7	0.76	6.8	0.09	1.9		II 2W	32,302,337,349,345 24
138	GGD 12-15	> 1	≥ 9	≥ 0.8	≥ 3	> 0.42	> 4.0		II W	1,49,7,337,338,345 25
139	S255-IRS1	25.8 ^h	87.9	1.7	50	0.06	1.8	0.18	II W	1,7,117,337,345
	S255									349
140	Mon R2E								H W	86,345
141	06114+1745	24 ^h	180	11	10	0.9	19	1.9	W	181,183,191,345
142	06291+0421	13 ^h	103	7	7.3	0.78	17	1.7	W	181,184,346
143	06308+0402	7	36.9	1.4	16	0.07	2.3		II W	3,341,346
144	AFGL961	20		80		11	70		2	1,7,50,286
		19	380	80	6	10.2	60			51,153
145	NGC2261							1.4E-3		74
	RMON									
146	Mon OB1H	1.6-2.3	17-53		1.2	1.1-8.2				47,53,147
147	Mon OB1D	16-30	92-520		6.9	0.36-11			II W	47,134,53,275
	NGC2264D									153,346
148	Mon OB1I	0.14-0.22	1.0-3.1		3.9	0.01-0.092				47,53
149	NGC2264G								2 I	206,291,335,336
150	NGC2264	1.6-7.2	11-150		1.1	0.32-25			WI	47,53,1,334,344
	Mon OB1C									147,349,346,345
151	Mon OB1G	1.3-1.9	26-93		1.8	1.9-21			2	47,53,132,291
152	S287N									86
	06453-0209									
153	BFS 56	0.3 ^h		0.14		0.011			W	86
	06567-0350									347
154	BiP 14	38 ^h		8.7		0.3			W	86
	06567-0355									347
155	CMa-W								W	86
	06568-1154									347
156	S287-B	0.3 ^h		0.12		0.003			H W	86,347
157	S287-C	0.3 ^h		0.07		0.003			H W	86
158	L1654								2W	86,287,347
159	S287-A	1.0 ^h		0.12		0.003			W	86,346
160	Z CMa ^c	0.82 ^h	3.2	0.145	3.1			3.62E-2	IIH	118,342

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot} \text{ km/s}$	E 10^{45} ergs	t 10^4 yr	L_m $L_{\odot}$	F $10^{-4} M_{\odot} \text{ km/s yr}$	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
161	CB54	0.75	4.4	0.26	2.8	0.077	1.6	0.15	I	178,179,231,329
162	07028-1100	44	267	13.5	24	0.5	11	1.1		421
163	L1660	3.2			7.8		1.8		IIH2W	91,280,255 338,339,346
164	07427-2400 G240.31+0.07	8.26 ^h	93	11	2.3	3.9	40		W	193,255,346
165	WB89 1099 HH120	1.07	1.97	6.0E-2	6.5	0.0076			H2 I	196,267,232
166	WB89 1135	151	419	16.1	37.9	0.36				196
167	HH46/47 ESO210-6A	0.03-0.3 ^h	1.9	2.2E-2-0.14	4	0.0045			H2	111,127,289
168	WB89 1173	28	58	1.8	15.8	0.094			2	196,285
169	WB89 1181	14	49	2	7.2	0.44			2	196,285
170	WB89 1187	55	171	7.7	16	0.4			2	196,285
171	WB89 1189	2.3	3.7	0.083	8	0.0086			2	196,285
172	08558-4347	13.1	107	10.4	5.0-26	1.29-4.41				256,259
173	WB89 1275	66 ^b	255	11.9	27.8	0.36			II	196,338,343
174	ϵ Ch I-C	0.15		0.014	2.7	4.5E-3				112
175	ϵ Ch I-N	0.21		0.037	3.8	8.1E-3				112
176	BHR71 IRS2	0.004							H	318
177	11590-6452 BHR71 IRS1	1.33 0.06	3.2-37.3	0.147-10.36	1.05	0.025-8.3			H2	177,188,288 318
178	12091-6129	115			0.8-27.3		26-8.61E5			208
179	12326-6245	320	5000	> 360	~ 1	100	1500		II W	219,338,343 416,349,417
180	12405-6238	243			1.2-37.6		39-1.288E6		II	208,338,343
181	12496-7650								H	130
182	12515-7641 HH52-53	2.4 ^h	7.5	0.25	8.1	0.033	1.3			130
183	12522-7640 HH54	0.68 ^{hb}	1.6	0.039	3.2	0.01	0.5		H	130,143
184	12553-7651 ^m	2.9	4.7	0.074	11	5.6E-3				357
185	13547-3944 CG12	0.05	1.6	0.017	2.7	4.2E-3	0.59			167
186	14562-6248	0.3	2.3	0.2					H	229
187	Circinus	50.6	540	85					H	229,186,226
	14563-6301	4.86	47.5	5.06	6.7					
188	14563-6250	0.3	2.3	0.2						229
189	14564-6258	0.6	6	0.6					H	229
190	14564-6254	4.02	38.8	4.17					H	229
191	14568-6304	0.15	1.2	0.11					H	229
192	14596-6320	0.4	4	0.4						229
193	14580-6303	0.4	4	0.4						229
194	14592-6311	0.4	4	0.4	1.4				H	229
195	16019-4903	90			0.9-28.2		18-5.90E5			208
196	16191-1936						0.004 $\pm$ 0.0003		II	190,342
197	L1686 ^b	> 0.003	> 0.016	> 0.09E-2	6.8	> 1.1E-4	> 2.3E-3	> 2.3E-4	H	28,31
198	Gss30 EL21	0.03			6.7				H2W	56,139,140,291 347
199	ρ Oph AS	4.72E-3	(3.63-7.0)E-2	(2.74-10.6)E-4	0.54-0.2 6	(3.73-30.0)E-3				378
200	ρ Oph A VLA1623	0.02	1	0.5	0.22	1.9	4.5	0.23	H2WI	106,232,347 291,293

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot} \text{ km/s}$	E 10^{45} ergs	t 10^4 yr	L_m $L_{\odot}$	F $10^{-4} M_{\odot} \text{ km/s yr}$	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
201	ρ Oph AN	1.34E-2 ^c	9.92E-2-	7.47E-3-	0.66-	9.15E-3-	0.084			378
		10.3E-3	2.88E-1	6.23E-2	0.2	2.495E-1				378
			5.2E-2-	2.8E-3-	1.23-	1.92E-3-				
			1.35E-1	1.75E-2	0.46	3.09E-2				
202	WL12						0.06 $\pm$ 0.006		H	190
203	EL29						0.15 $\pm$ 0.014			190
204	ρ Oph-B	0.17 ^h								92
205	IRS43						0.15 $\pm$ 0.012		W	190,345
206	ρ Oph B2	1.66E-2 ^c	7.4E-2-	3.4E-3-	5.6-	8.1E-4-	0.094			378
			2.06E-1	2.5E-2	0.74	4.6E-2				
207	L1681B	0.03 ^h			0.4		0.4		II WI	84,345,375
208	IRS48						0.30 $\pm$ 0.061			190
209	IRS51						0.055 $\pm$ 0.018		W	190,345
210	L1709A ^d									31
211	L1709									86
212	ρ Oph-S									86,234,331,233
213	Re 13	1.25				0.076	1		H	114,115
214	V346N or FuOri	0.37 ^{bc}	3.42	0.4	1		3.42	0.133	H2	114,118,115,290
215	ρ Oph-E	0.59 ^h	3.7	0.27	9	0.0183			II WI	33,131,105,231
	L1689N	0.8	12.8	0.05	1.7-5.8	2.1	16.3	> 0.1		136,233,234,33
		0.15	0.7		0.9	0.05	0.7			131,369,339
	16293-2422	0.39			2.5	0.43	3.2			228,345,105
216	16295-4452									320
217	16306-4455									189
218	16311-4457	1.36				0.114	1.5			189
219	L43	0.11	0.12	0.0052	17	2E-4	0.005	0.5E-3	II 2W	22,23,69,215,13
	RNO91	0.04								383,347,358,374
220	16442-0930								W	86,84,23,347
	L255									
221	L146									86
222	17081-2712	0.342	1.44	0.0629	4	0.0099				257
	or B59-MMS1									
223	L100									93,403
224	NGC6334	86 ^h	502	41	1.3	3.31			W	88,57,59,349,353
225	NGC6334B	2344			2.82	6.2	141		II W	239,338
										120,349,352
226	NGC6334I ^c	2.3	92	43	0.23	170			II 2W	96,291,255
		3317			0.6	71.9	766			239,349,352
227	CB81									178,179
228	17364-1946								II	168,342
	L219									
229	W28-A2	40.4	593	113				56.4	II W	152,338,342
	G05.89-0.4	70	1600	800	0.2	1600				94,349,345
										255
230	H 36								II	322,338
231	M8E	9.6				2.9	45		II	7,1,342
			39.2	18				50		58,348
232	G9.62+0.19F ^m	$\sim$ 50			0.8				II	255,355
233	18128-1943	14.1	67	3.7	14	0.14-0.36				260
234	18134-1942	37	180	10.2	13	0.36-2.24			W	260,346
235	18139-1952	180	610	26	7	1.3-11				260
236	L483								WI	124,231,93
										232,329,347
237	18150-2016	3.2	11.6	0.44	24	0.03-0.09				260
238	18151-1208	12 ^h	106	9	7.1	1.1	15		W	237,347
239	GGD27	460 ^h	5300	78	10	6	5300		IIH2W	15,294,337,338,339
		407 ^h			7.63	15.8	147			239,345
240	18182-1433	203	1665	150	15.3	8.2	110		W	237,349,345
		30	246	23	8.9	2.1	28	34		

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot} \text{ km/s}$	E 10^{45} ergs	t 10^4 yr	L_m $L_{\odot}$	F $10^{-4} M_{\odot} \text{ km/s yr}$	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
241	18196-1331 CRL2136	50			~ 10				II 2W	164,295,340,342 347
242	18258-0737	34.5 ^c	128	15.4	6.27	2.1	3.7		II	203,339
243	18264-1152	405 ^h 31 ^h	6818 534	1100 90	7 3.7	135 20	980 140		W	237,314 237
244	L379IRAS(3)	1.8 21.3 ^h			≤ 2.4 3.3	~ 0.35 32	5.6 30-160		II W	42,339 116,347 327
	L379IRS1	0.67	8.7	2.6						
245	Serpens SMM1-11 SMM1-4,8.9?	~ 1.8 0.3	12.3 2	18.1 3					H WI	331,345,415 365,406
246	S68 Firs1 SVS 4								IIH2WI ^a	1,78,7,292,331,361,76 405
247	L379IRAS(2)	10.7	92.4		6.6		14		II W	91,42,342,346
248	64 ^b ,W4	4.1 ^h	11.1	0.28	24	0.009	0.45	4.5E-3		173,420
249	18316-0602								II 2W	303,255,337,339
	G25.65+1.05	7.64 ^h	136	27	1.8	12.4	76			193,346
250	18331-0035								H	395,396
251	18345-0641	143 ^h	1841	240	15	13.1	120		II	237,340
252	18470-0044	102 ^h	1050	110	14.8	6	71			237
253	18537+0749	14.0 ^c	240	5.5	2.3	33.3	1.21		II W	203,341,349,346,345
254	G35.2-0.74	~ 60 73 ^h		800	2 4.6	320 5.1	1.00E+03 2.70E+02	8	II W	72,73,54,191 228,337,340,349,345
255	18566+0408	32 ^h	540	91	7.5	10	72		W	237,347
256	R CrA	0.13			2.7			1.5E-2	H2	8,293
257	19012+0536	134 ^h 38 ^h	1339 383	130 38	12.2 13	8.9 2.4	110 29			237 237
258	19035+0641	3 ^h	28	3	2.9	0.9	10		W	237,349,345
259	W49	600 ^h	3500	670					II W	95,341,340,349,345
260	G45.07+0.13	$10^{1.65c}$	$10^{2.7}$		4	$10^{0.74}$	$10^{1.9}$		II W	202,255,340,341,349,345
261	G45.12+0.13	$10^{3.4}$	$10^{4.38}$		18.6	$10^{2.07}$	$10^{3.12}$		II W	202,255,340,341,349,345
262	L723 2 Flows	0.2-0.5 0.38 4.7 0.42	1.0-5.0 29.5	0.07-0.53	4.7-3 3.4 5	0.01-0.1 0.044 0.4	0.2-1.6 0.5	0.02-0.16	H2WI	14,7,153,241,232,418 228,244 154 383
263	CB188	5.1	2	0.08	2.6	0.024	0.7	0.066		240,178,179
264	L673 SMM1	0.0052		$7.4E-4$	3.3		0.057			422,423
265	L673	1.6	3.9	0.013	9	0.012				61
266	AS353	> 0.05	≥ 0.38	≥ 0.01	1.4	> 0.003		$> 7E-3$	2	62,7,8,311
267	19213+1723	3.9	24.8	1.9					II W	207,341,346
268	19217+1651	108 ^h	1961	360	7.5	38.8	260			237
269	L778	0.08	0.28	$2.1E-2$	8	$1.1E-3$	0.019	$1.9E-3$	II	22,13,28,375
270	19266+1745	35 ^h 0.04 ^h	311 0.3	28 0.03	18.4 1	1.3 0.02	17 0.3			237 237
271	19282+1814	0.86	7.1	$2.5E-7$	6	0.3	2.4	0.12	II	419,337,341
272	B335	0.1-0.8 0.44	0.3-4.8	0.01-0.24	9.4-6.0 11	0.009-0.03 0.019	0.03-0.8 0.36	0.3-7.5E-2	H2 I	14,30,230,234,331 228,298,231,232
273	19374+2352	17.1 ^c	122	10.8				10.2	II W	207,341,346
274	1548C27	0.18	~ 1.0					$1E-2$	H	384, 385
275	19410+2336(L)	80 ^c 13.5 ^c	622 113.6	4.8 7.9	8 3.2	4.9 2.03	7.9 3.2		WI	203,349,345,424 203
		1423 ^h	31896	7100	5.9	982.9	5400			237
		153 ^h	3434	770	3.8	165.3	900			237
276	19410+2336(U)	49 ^c 8.2 ^c	336 56.2	230 3.8	8 3.6	2.2 0.89	3.8 15.5		W	203,349 345
		173 ^h 18 ^h	3766 405	820 88	6.2 4	108.3 18.2	610 100			237 237
277	19411+2306	46 ^h 12 ^h	649 162	90 23	6.9 6.9	10.8 2.7	94 24			237 237
278	L810	1.87-28	6.96-146	0.282-8.39	10	0.02-0.6	0.001-0.0209	0.07-1	W	97
	CB205	0.64	5.5	0.46	1.5	0.26	3.3	0.33		240,346
279	S87	34	102	9	5.4	1.4	19	1.8	II W	1,7,125,337,341
280	S88		48-480	3-30	7				W	16,1,77,349
	S88 B	867			4.68	222	1030		II	239,341,345

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot}\text{km/s}$	E 10^{45}ergs	t 10^4yr	L_m $L_{\odot}$	F $10^{-4}M_{\odot}\text{km/s yr}$	$\dot{M}^o$ $10^{-5}M_{\odot}/\text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
281	19471+2641	250c	1653	108	4.5	2.3	42		II W	203,347
		67.5	453.1	6.7	21	1.2	5.1			203,341,347
282	19520+2759	5-13	42-113	8-20	< 10	>0.5-1.4	> 4		W	87,346
283	19529+2704	24.91	290.74	28	5.2	4.4	56	5.6	II	198,341
		35.8	417.84	40	6.2	5.3	67	6.7		198
284	19550+3248	1.45			9.4	0.046		1.48	W	119
		5.77			3.67	0.35	2.9			239
285	K3-50								II W	16,337,341,349,345
	19598+3324									
286	CB214	0.42	2.2	0.11	2.7	0.033	0.7	0.066		240,178,179,345
287	CB216	0.49	2.8	0.18	2.3	0.062	1.1	0.11		240,178,179
288	20050+2720	1.7	25.4	7.65	2.2	~ 16	~ 50	~ 20	WI	248,210,232,231,331,346
289	20056+3350	2.3	20.6	2.8		~ 20		~ 50	W	207,346
290	CB217	0.14	0.4	0.018	2.6	0.007	0.2	0.022		240,178,179
291	20062+3350								W	207,346
292	20106+3545	1.9	6.7	0.3						207
293	20126+4104	67 ^h	2100	670	2.1	260	1.00E+03		II 2W	116,299,255,363
		3.8 ^c	230	220	0.094	300	2.10E+03			250,346
		53	403	51						207
294	20188+3928								W	63,123,346
		0.3	2.2	0.2	4.54	1.1	26.6		II	239,339
295	G75N	19 ^h	260	40	5.3	6.2	40	36	W	225,349
296	G75C	58 ^h	1060	200	3.7	45	28	160	II W	225,341,349,345 77
297	G75E	20.7 ^h	350	63	4.7	11	74	43	II W	194,225,303,341,349,345
298	G75.78NE	72	970	130	3	36	320	2	II	194,255
299	20216+4107	6 ^h	43	3	3.5	0.8	13		W	237,141,347
300	V1515Cyg									118,137
301	20227+4154 ^s								W	123,346
302	20228+4215 ^s								II	123,339
303	20231+3440 SMM1	2.1 ^c	18.7	1.8	5.6	0.34				372
304	20231+3440 SMM2	0.15 ^c	1.08	0.08	4	0.016				372
305	20272+4021 ^s									123
306	AFGL2591	33.4	269	23				6.8	IIH2W	152,349,345,348
	20275+4001									1,64,123,298,255
307	20281+4006 ^s									123
308	20286+4105	16.7	45.6	1.6					II	207,339,346
309	20293+3952#1	9 ^h	270	78	1.1	59.2	250		W	237,314
		4 ^h	114	33	1.3	20.6	87			237
310	20293+3952#2	1.1 ^h	21	6	0.5	9.1	42			237
		0.4 ^h	8	2	0.6	3.2	15			237
311	20343+4129#1	2 ^h	9	0.4	3.8	0.1	2			237
312	20343+4129#2	0.4 ^h	1	0.07	4.2	0.01	0.4			237
313	20353+6742						0.047 $\pm$ 0.010		H	190
	L1152									
314	W75-N	130 ^h	955	100	2.3	0.99			II 2W	88,48,302
		548			1.93	46.4	352			239,349
		48	~ 700	110	≥ 4.4	22				255,247,345
315	DR21	170 ^h	1203	140	2	9.6			II 2WI	88,65,297,337,424
		3.30E+03	2.40E+04	2.30E+03						148,255,1,349,345,362
316	L1036									86
317	L1157	0.18 ^h	1.1	0.076	2.4	0.03			H2WI	122,86,291,331,231
										214,232,347,399
318	PVcep	1.7 ^h				0.03	8.5		H W	7,345
		0.8-2.6			5-15			0.04-0.2		66
319	20520+6003						0.25 $\pm$ 0.029			190
	L1082A									
320	20526+5958						0.028 $\pm$ 0.003			190
	L1082B									

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot} \text{ km/s}$	E 10^{45} ergs	t 10^4 yr	L_m $L_{\odot}$	F $10^{-4} M_{\odot} \text{ km/s yr}$	$\dot{M}^o$ $10^{-5} M_{\odot} / \text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
321	V1057Cyg	0.09			6			2.4E-3	W	8,12
322	20582+7724	0.06			2	0.04			IIH2	71,301,400,401
323	V1331Cyg	0.6			7			2.0E-2	H	8
324	L988-a			1.3	5	0.1			W	98,346
325	21015+6757	1.3	6.4	0.28	24	0.009	0.27	8.9E-3		397,398
326	L1172D	0.88	2.3	0.14	17	3.6E-3	0.066	6.6E-3	I	22,13,28,334
327	L988-e			1.4	4.5	0.2				98
328	L988-f			3.5	3.3	0.4		> 0.024		98
329	B361 ^b	> 0.025	> 0.055	> 0.0013	2.4	> 4.3E-4	> 2.4E-2	> 2.3E-3	W	31,28,347
330	L944 SMM1	0.059		6.40E-03	3.3		5.7			422,423
331	CB230	0.46	1.7	0.07	2.6	2.2-2	0.7	6.6E-2		240,178,179
332	IC1396-W									86
333	21307+5049	19.3	92.1	7.2					II W	207,341,346
334	21334+5039								II 2W	113,339,346
335	CB232	0.44	2.2	0.11	3.4	2.6E-2	0.7	6.6E-2		178,179,240
336	V645Cyg	56.1 ^h			50			0.14	IIH W	8,70,345
										1,7,349
337	GN21.38.9	0.07			30	7.3E-5		1E-3	IIH W	150,254,326,346
338	IC1396-N BIMA1	1.2	0.2	0.04			0.7			373
339	IC1396-N ⁿ	0.3 ^h	1.2	0.05	3	0.02			H2W	89,254,300,328
		0.75 ^h	16	3.5	1.4	1.2-2.0	7-11			116,346
340	IC1396-N BIMA2	13.6	3.6	1.2			13.6		W	373,393,414
341	1413+5442 or G98.04	40	240	15	9.1	1.4	27		II W	194,255,350,346
342	NGC7129 FIR ^u	13.9	49	2	10	$10^{-0.8}$	5		H2W	7,18,78,266,345
343	NGC7129 FIRS2-MM1				> 0.3			0.3	H2W	317,266,345,374
344	NGC7129 FIRS2-IR				> 0.5				H2W	317,266,345,374
345	NGC7129 FIRS1-IRS6				> 0.8				H2W	317,266,345
346	NGC7129 FIRS1-MM1									317,345
347	LKH α 234	23.5	84	3	10	$10^{-0.6}$	7.9	0.4	H2W	18,78,266,304
						1	20			7,345
348	21428+4732	0.09	0.33	0.014	4.6					258
349	21429+4726	0.1							H	162
		0.57	2.58	0.12	6.9					258
350	21432+4719	2.04							H W	162,345
		0.04	0.18	0.01	22					258
351	21441+4722	0.206								162
		0.21	1.03	0.05	11					258
352	IC1396-E									86,254
353	EL1-12	0.93			3			0.25		52,28,22,7
	L1031B	1.2								153
	IC5416									137
		2.56	15.9	1.15	4.1					258
354	21461+4722	0.05								162
		0.15	0.36	0.01	5.7					258
355	BBD+46 ^o 3471	1.4 ^h			7			0.04		8
356	21519+5613	65.6	387.9	28.6					W	207
357	22051+5848									168,91
	L1165									
358	22103+5828	3.3	18.4	1.08	3.5	0.2				185
359	22134+5834	17 ^h	242	34	3.5	7.9	69		II 2W	237,306,321,374
		19	124		15					180,345
360	22142+5206	33	219	15.7	23				W	187,345

Table 1b Parameters of CO molecular outflows(continued)

No.	Name	M $M_{\odot}$	P $M_{\odot}\text{km/s}$	E 10^{45}ergs	t 10^4yr	L_m $L_{\odot}$	F $10^{-4}M_{\odot}\text{km/s yr}$	$\dot{M}^o$ $10^{-5}M_{\odot}/\text{yr}$	IIH2WI ^a	Ref.
(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)
361	22172+5549	9	37.1	2.1					II W	207,345,374
362	S140	64				90	600		IIH2W	1,7,332,255,337
		24.4	173	15	1.5	7.8	120			11,349,332
		6	24	1.2						82,345
		25.24	162.2	12.42				1.0-5.2		79,151,152
		24.4			2.6	8.4	53			228
		24.4	488	98	11	8.4		94		255
363	S140-N								H2	33,36,264,332
	22175+6317									
364	L1204-A	0.5	3.2	0.2					W	207,346
	22198+6336									86
365	L1204-B								H W	86,332
	22199+6322									346
366	L1221	1.3	2.0-9.6	0.033-	5	0.0056-			IIH W	86,149,374
	22266+6845			0.71		0.12				184,418
		0.34								383
367	L1203								W	86
	22267+6244									347
368	L1206	0.2 ^h	0.9	0.04	2	0.01			H	89,328
369	22305+5803	15.6	79.4	4.9					W	207,345
370	22336+6855				2.5					176
371	22343+7501	1.95	11.7	0.69	13.6	0.042			H W	68,171,347
372	22376+7455	0.116	1.14	0.111	2.3	0.039			H WI	68,171,231,235,334
	L1251									235,347
373	L1211								W	86
	22453+6146									347
374	22475+5939	59.9	633.4	51	29	1.4	22	2.2	W	199,349,345
	S146	51	538	43	5.4	6.4	99	0.5	II	191,203,341
375	22506+5944	55	670	69	2.3	24	290	5.8	W	191
	10860+0049									345
376	Cep A	10		62		25	120		IIH2WI	1,7,60,246,255,337,341
		8.8	85.7	10.02	0.2			2.8		152,305,424
		> 10			> 0.8					55,345
		5			10			0.2		8
		5	100	20	6					116,349
377	22570+5912	73 ^h	911	110	12.1	7.8	75		W	237,346
378	23011+6126	0.21	1.74		< 4				H W	86,204,389
	Cep E									345
379	23032+5937	34.2 ^h	172.4	9.74	10				W	99,345
380	23033+5951	32 ^h	566	100	5	16.9	110		2W	237,306,345
381	Cep C								W	86,345
	23037+6213									
382	MBM 55	4.1	8.2	0.162	22	0.0059		3.6		100
383	NGC7538 ^f	45	275	20			760	1	II 2W	1,44,95,308,341,349,345
384	NGC7538	16	251	40	2.4	15.7	1.10E+02	10	II 2WI	76,302,349,424
	IRS11									345
385	NGC7538	16	302	60	2.4	23.5	1.40E+02	10	II 2W	76,308,349
	IRS1									345
386	NGC7538 A ^d	11	251	50	3	13.1	7.90E+01	8	W	76,349,345
387	NGC7538	34	503	100	4.1	26.1	1.60E+02	20	W	76,349,348
	IRS9	49.5	410	40	1.6	40		320		312,345
388	23139+5939	15.5 ^h	82.5	4.84	7				II W	99,345
	G111.25	16	110	8.4	3.7	1.8	31			194,255
		57 ^h	662	77	5.4	11.7	120			237
389	23140+6121	3.5	34.5	2.7					W	207,347
390	23151+5912	21 ^h	597	180	2	72.7	290		W	237,350
		8.9 ^h	47.3	3	4.4					99,345
391	MWC1080	100 ^h	550	35.8		0.96	16	26	IIH I	2,7,375
										151
392	L1246 SMM1	0.019		5.9E-4	4.6		0.72			422,423
393	CB244	0.07 ^h -0.1 ^h		0.001-0.1	0.9			3	I	84,93,179,334
	L1262	0.4	2	0.1	2.7	0.031	0.7	0.066		240,231
394	23314+6033	5	17.7	0.7					II W	207,341,345
395	23385+6053	23	226	17	4	3.5	56	1.1	II W	191,359,341
		45.7	188.1	8.7						207,345
396	23545+6508	3.5	31.7	2.4	12	0.15	2.5	0.25	II W	199,339,347,374
397	CB247 ^b									178,179,240

A study of high velocity molecular outflows with an up-to-date sample [★]

Yuefang Wu¹, Yue Wei¹, Ming Zhao¹, Yong Shi¹, Wentao Yu¹, Shengli Qin¹³, and Maohai Huang²³

¹ Astronomy Department, CAS-PKU Joint Beijing Astrophysics Center, Peking University, Beijing 100871, China

² Osservatorio Astronomico di Trieste, via Tiepolo n. 11, Trieste 34131, Italy

³ National Astronomy Observatory of China, Beijing 100012, China

Received ?, ?; accepted ?, ?

Abstract. A statistical study of the properties of molecular outflows is performed based on an up-to-date sample. 391 outflows were identified in published articles or preprints before February 28, 2003. The parameters of position, morphology, mass, energy, outflow dynamics and central source luminosity are presented for each outflow source. Outflow lobe polarity is known for all the sources, and 84% are found to be bipolar. The sources are divided into low mass and high mass groups according to either the available bolometric luminosity of the central source or the outflow mass. The pace of discovery of outflows over the past seven years has increased much more rapidly than in previous periods. Surveys for outflows are still continuing. The number of high-mass outflows detected (139) has considerably increased, showing that they are commonly associated with massive as well as low mass stars. Energetic mass ejection may be a common aspect of the formation of high mass as well as low mass stars. Outflow masses are correlated strongly with bolometric luminosity of the center sources, which was obtained for the first time. There are also correlations between the central source luminosity and the parameters of mechanical luminosity and the thrust or force necessary to drive the outflow. The results show that flow mass, momentum and energy depend on the nature of the central source. Despite their similarity, there are differences between the high mass and low mass outflows. Low mass outflows are more collimated than high mass outflows. On average, the mass of high mass sources can be more than two orders of magnitude larger than those of low mass outflows. The relation between flow mass and dynamical time appears to differ for the two types of outflows. Low mass sources make up 90% of outflows associated with HH objects while high mass outflows make up 61% of the sources associated with H₂O masers. Sources with characteristics of collapse or infall comprise 12% of the entire outflow sample. The spatial distribution of the outflow sources in the Galaxy is presented and the local occurrence rate is compared with the stellar birth rate.

Key words. star: formation – stars: winds – ISM: jets and outflows – ISM: kinematics and dynamics

1. Introduction

As one of the most exciting discoveries in astronomy in the past three decades, high velocity molecular outflows continue to attract great attention from researchers. Since their first discovery in 1976 (Zuckerman et al. 1976; Kwan & Scoville 1976; Zuckerman & Palmer 1975), outflows have been detected at a high rate. 67 had been dis-

covered by 1984 (Lada 1985), 144 by 1989 (Fukui 1989), and 264 by 1996 (Wu et al. 1996).

Outflows were thought to be the earliest observable signatures of star formation. Although evidence of mass loss for young stars had been detected previously in the optical spectrum (Herbig 1960; Kuhl 1964), a better understanding of such mass loss in early evolutionary phases was only acquired after the discovery of molecular outflows in the material surrounding the embedded young stellar objects (YSOs). Molecular outflows show various characteristics such as radio and optical jets, atomic hydrogen stellar wind, molecular hydrogen jets and interstellar masers. The stars eject mass in energetic flows dur-

Send offprint requests to: Yuefang Wu, e-mail: yfwu@bac.pku.edu.cn

[★] Table 1 is only available in electronic at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.125.5) or via <http://cdsweb.u-strasbg.fr/Abstract.html>

ing formation (Lada 1985; Lizano et al. 1988; Natta et al. 1988; Lane 1989; Felli et al. 1992; Russell et al. 1992; Matzner & McKee 1999; Bally et al. 2001; Reipurth et al. 2002). In the 1990s, progress was made in observing and understanding the collapse of protostars (Zhou 1992; Evans 1999; Myers et al. 1996). According to theoretical models, collapses should occur before outflow (Shu et al. 1987). However, nearly all the candidates of collapsing protostars are found in outflow sources.

When advancement of observational techniques allowed the detection of high velocity molecular outflows by high J transitions, extremely high-velocity molecular outflows or highly collimated jet-like outflows were studied extensively (Koo 1989; Bachiller et al. 1991; Choi et al. 1993; Narayanan & Walker 1996).

More recently, considerable progress in outflow detection has been made in distant and complex high-mass star formation regions (Beuther et al. 2002c; Zhang et al. 2001; Shepherd & Churchwell 1996b; Wu et al. 1999). In the surveys of high-mass star formation regions conducted by Beuther et al. (2002c) and Zhang et al. (2001), the outflow detection rates reached 80% and 90%, respectively. Recent studies show that the sources in these surveys are either precursors of ultracompact (UC) HII regions, or are at very early evolutionary stages (Molinari et al. 2002; Beuther et al. 2002b).

The outflow parameters and their correlations have been investigated (Rodríguez et al. 1982; Bally & Lada 1983; Cabrit & Bertout 1992; Shepherd & Churchwell 1996a; Ridge & Moore 2001). Mechanisms of outflow collimation and the forces driving the outflows have been proposed as well. However, the correlation between morphology, physical parameters and the central source conditions are still not well understood. For example, it is not clear if the collimation is correlated with the driving force. The correlation among bolometric luminosity of the center source, the required force and mechanical luminosity was investigated by Bally & Lada (1983) and Rodríguez et al. (1982). As the correlation was discovered to be weak, a larger sample is required.

To further understand the mechanisms of high velocity outflows, we conducted an analysis based on an up-to-date sample. Our calculations and conclusions are presented in this paper.

The used data in this paper are heterogeneous. The outflows were detected at different stages of the young stellar objects: high mass sources from pre-UCHII region to UCHII region and low mass sources from Class 0 to FU Orionis. The observational sensitivity and the calculation method also varied from source to source. However, we believe that our sample is sufficiently large to provide the overall physical characteristics of outflows. Sect. 2 presents the sample, the catalogue and a brief analysis. The morphology analysis in Sect. 3. Sect. 4 discusses the physical

parameters. The associated objects or phenomena of outflows are discussed in Sect. 5. Sect. 6 describes the spatial distribution of outflows. We give a summary in Sect. 7.

2. The catalogue

2.1. The sample

We compiled outflow sources mapped mainly in emission lines of low transitions ($J = 1 - 0$ and $J = 2 - 1$) of CO showing evidence of large scale red and blue lobes. The catalogue of 1996 (Wu et al. 1996) included 28 sources without CO maps (with superscript a in the source name). We exclude those sources that have not been mapped so far. If an outflow was detected with CO $J = 3 - 2$ or higher transition lines, it is noted in the table accordingly. Some authors detected molecular outflows with other molecular species, such as SiO, CS, HCO^+ (e.g. Megeath & Tieftrunk 1999; Wolf-Chase et al. 1998; Garay et al. 1998; Hofner et al. 2001). These species are not included in this catalogue.

The sources presented in this new catalogue were identified as high velocity molecular outflows in published findings or in preprints published before February 28, 2003. There are 397 sources; 391 show single outflows, 6 were observed as single outflows but were shown later to be of multiple sources. These six were presented in the catalogue, but are not included in our analysis.

2.2. The tables

Table 1a and 1b list the high velocity molecular outflows and their parameters. Some of the outflows have been investigated by multiple authors, and have been mapped more than once. For these sources several entries are listed. Each entry corresponds to the references listed in the last column of Table 1b. There are 24 columns for each entry. Column 1-13 are in Table 1a, and the remaining columns are in Table 1b. The sources are presented in order of right ascension.

Table 1a contains the basic parameters of the outflows:

- Column 1 (No.) is the sequence number of the outflow.
- Column 2 (Name) the source name and the alternative name if any.
- Column 3 (α) and 4 (δ) are the right ascension and declination (1950.0).
- Column 5 (l) and 6 (b) are the galactic longitude and latitude.
- Column 7 (D) and 8 (Z) are the distance from the sun and altitude from the Galactic plane.
- Column 9 (ΔV) lists the full line-width at 0.1-0.2K above the baseline. For extremely high velocity sources, an additional line-width measured at tens of milli-kelvins is provided.

- Column 10 (Po.) shows the polarity of the outflow. Here, ‘Bi’ represents bipolar outflow; ‘MB’ and ‘MR’ indicate blue and red monopolar outflows, ‘Multi’ means multiple outflows and ‘Iso’ isotropic outflows.
- Column 11 ($R_{\max}$) and 12 (R_{coll}) present the size and the collimation factor, respectively. To calculate the collimation factor, we first consider an outflow as an ellipsoid. Then we measure the area and maximum angular extent (the major radius) of the outflow, and calculate the minor radius. The ratio of the major and minor radii is the collimation factor (Bally & Lada 1983). The value underlined in column 12 is quoted from references where the same method was employed.
- Column 13 (L_{bol}) shows the bolometric luminosity of the center source which is the deriving source or its candidate, identified by infrared photometry or by IRAS data.

The physical parameters are listed in Table 1b (column 14-24), mainly derived from lower level CO lines (CO $J = 1-0$, $2-1$). If the parameters were calculated with CO $J = 3-2$ or higher levels, we note them with ‘c’. For extremely high velocity outflows we only list the total velocity range.

- Columns 14 and 15 contain source numbers and their names corresponding to Table 1a column 1 and 2.
- Column 16 (M) is the outflow mass calculated by integrating CO emission from the entire area of the wings (Goldsmith et al. 1984; Snell et al. 1984; Lada 1985). The mass obtained under optically thick conditions are noted ‘h’. The rest are calculated in the optically thin regime.
- The momentum (P), kinetic energy (E) and dynamic time (t) of the outflows are listed in column 17, 18 and 19, respectively (Lada 1985).
- The mechanical luminosity (L_m) of the outflow and force (F) derived from outflow physical parameters (Bally & Lada 1983; Goldsmith et al. 1984; Snell et al. 1984) are shown in column 20 and 21, respectively.
- Column 22 ($\dot{M}$) indicates the mass loss rate of the center stellar source, which was calculated with the outflow parameters and terminal velocity of the stellar wind (Bally & Lada 1983; Goldsmith et al. 1984; Snell et al. 1984; Levreault 1988) and the corresponding reference of the outflow.
- Column 23 (IIH2WI) presents the phenomena associated with the outflows. ‘II’ stands for UCHII regions, ‘H’ for optical jet or HH objects, ‘2’ for H₂ jet, ‘W’ for water maser, and ‘I’ for infall or collapse candidates.
- Column 24 (Ref.) presents the references.

2.3. Classification

The bolometric luminosity of the center source of the catalogued outflows ranges from one tenth to more than 10^6 times solar luminosity ($L_{\odot}$). We adopt $L_{\text{bol}} = 10^3 L_{\odot}$ as a criterion to distinguish between high-mass sources and low-mass sources. Young high-mass stellar sources are brighter than $10^3 L_{\odot}$ at minimum. On the H-R diagram, L_{bol} of low-mass and intermediate mass sources is mostly below $10^3 L_{\odot}$ (Levreault 1985, and references therein). The criterion of $10^3 L_{\odot}$ was similarly employed in the comparative study of different mass molecular outflows or cores (Shepherd & Churchwell 1996a; Fukui 1989; Wu et al. 2001).

Many sources have more than one set of parameters derived from different observations, therefore we classify them according to the following considerations. First we only classified sources whose multiple measurements of masses have consistent L_{bol} values either larger or smaller than $1000 L_{\odot}$. In the classified sources, 292 sources have available L_{bol} measurements, of which 120 (41 %) have $L_{\text{bol}} > 10^3 L_{\odot}$, and 172 (59 %) have $L_{\text{bol}} \leq 10^3 L_{\odot}$. The rest of the sources have no available L_{bol} , but some have available outflow gas masses. Of the 113 sources with $L_{\text{bol}} > 10^3 L_{\odot}$ and with available masses, 94 (82%) have $M > 3 M_{\odot}$; for 138 sources with $L_{\text{bol}} \leq 10^3 L_{\odot}$ and with available masses, 111 (80%) have $M \leq 3 M_{\odot}$. Sources whose masses observed by different authors have values both larger or smaller than $3 M_{\odot}$ were not taken into account in the above calculation. Thus, the two percentages cited above are lower limits. Further statistics show that the outflow mass is correlated with L_{bol} (see Sect. 4.1). For sources without L_{bol} but with mass available, we use $3 M_{\odot}$ as a criterion to classify them. Among these sources there are 19 with $M > 3 M_{\odot}$ and 51 with $M \leq 3 M_{\odot}$. We refer to the 139 sources with $L_{\text{bol}} > 10^3 L_{\odot}$ (120) or $M > 3 M_{\odot}$ (19) as the high mass group and the 223 sources with $L_{\text{bol}} \leq 10^3 L_{\odot}$ (172) or $M \leq 3 M_{\odot}$ (51) as the low mass group in the following text. Currently we have not found the mass or bolometric luminosity for 29 sources. These sources need further examination.

2.4. Development of outflow detections

New outflows have been detected and surveyed at an accelerated rate in the past two decades. There are now 397 identified sources (including 6 unresolved outflows). The number of sources is seven times greater than that catalogued in 1985 (which had at most 55 mapped) (Lada 1985), 244 more than that of 1989 (Fukui 1989), and 159 more than that of 1996 (Wu et al. 1996).

Fig. 1a is the diagram of the number of outflows mapped in four periods, which is defined with the available catalogues of outflows. Period I is from 1976 to 1984 with

55 detected outflows (Lada 1985); Period II is from 1985 to 1989, with 95 outflows (Fukui 1989); Period III is from 1990 to 1995, with 88 outflows (Wu et al. 1996); Period IV is from 1996 to 2003 with 159 outflows (this work). The number of detected outflows has increased, especially in period IV (see Fig. 1a). Fig. 1b is the average number of mapped outflows in these four periods. The average number of outflows per year are 6.9, 19, 14.6 and 22.7 for the four periods respectively. The trend of increasing number of detected outflows is evident, showing the development of outflow detection in recent years.

In period II mainly low mass outflows were added (Fukui 1989), while in period IV the number of high mass sources increased rapidly. Among the sources detected up to 1995, high mass sources occupy only 31% (65 of 210 outflow sources in the high mass group with the same classification). Now there are 139 in the high mass group among 362 sources with available L_{bol} or M . The ratio is 38%. The results show that high velocity molecular outflows are common in high mass star formation regions considering the fewer samples of high mass stars. It may suggest that high mass stars still form through an accretion-outflow dynamic process, like the formation process of low mass stars.

Is the percentage of high mass here beyond that expected from the IMF? Here 38% is the ratio of sources in the ‘high mass group’ to the total number of sources in both the high and low mass groups. Part of the ‘high mass group’ are actually intermediate mass sources. Considering the sources with $L_{\text{bol}} > 10^4 L_{\odot}$, the percentage drops to 23% (83/362). The percentage of sources associated with UCHII regions (23% (90/391)) is also lower than 38%. This percentage still exceeds what is expected from the IMF. But observed IMFs are derived from counting main-sequence stars in solar neighborhood (Kroupa et al. 1990; Salpeter 1955). According to Fig. 9, massive outflows are easier to detect at increasing distance. Outside the 2 kpc circle, most outflows belong to the high mass group, while within 1 kpc, there are only 10% (16/167) high mass outflows.

A notable trend in outflow research is that interferometers are being applied to measure the outflow regions. Massive outflows have been observed with higher spatial resolution and sensitivity in regions such as IRAS 05358+3543, G196.16 and NGC 7129 (Beuther et al. 2002a; Shepherd et al. 1998; Fuente et al. 2001).

3. Morphology and collimation

The polarity was known for all the outflows. 327 (84%) sources show bipolar structure; 50 (13%) sources are mono-polar, of which 28 are red polar. There are 12 (3%) multi-polar sources; if a source was detected as bipolar first but was later found to have multiple polarities we

count it as multi-polar. Two sources are still isotropic (No. 231 M8E and No. 391 MWC 1080). Compared with the sources cataloged before (Wu et al. 1996; Lada 1985), the number of multi-polar sources has increased (there is only one in the previous periods) and the number of isotropic sources decreased (4 in Period I, 3 in Period III), which can be attributed to the enhanced spatial resolution of the detection equipment.

The collimation factor is used to investigate the morphology of outflows. Different methods were employed to analyze collimation properties of outflows (Cabrit & Bertout 1992; Levreault 1985). In this paper the collimation factor was obtained according to Bally & Lada (1983). Here we analyze the relationship between the collimation factor, the bolometric luminosity of the central source and the angle sizes.

As shown in Table 1a, there are 213 sources with available collimation factors. The average value of the collimation factor for all the sources is 2.45, with a standard deviation of 1.74, which is almost the same as that catalogued in period I and III (Lada 1985; Wu & Huang 1998). The average collimation values for high mass and low mass group members are 2.05 and 2.81, with a standard deviation 0.96 and 2.16 respectively. The above values are similar to those in period III, 1.99 and 2.54 for these two groups according to the same statistical method (Wu & Huang 1998). These results show that the low mass outflows are better collimated than the high mass ones. Although the overall instrumental angular resolution in period IV has increased,

the observed collimation has not improved much as a whole. This is likely due to the increased number of massive outflows.

Figure 2 is the plot of R_{coll} vs. L_{bol} . Solid triangles indicate low mass sources and squares indicate high mass ones. The solid line is a linear fit of the plots: $\log(R_{\text{coll}}) = (0.44 \pm 0.03) + (-0.04 \pm 0.01) \log(L_{\text{bol}})$; the correlation coefficient is poor, $R = -0.29$. There is a slight difference between high mass and low mass outflow collimations. To analyze the effect of angle size on the collimation factor, we plot R_{coll} vs. the angular size in Fig. 3. It shows that there is no obvious correlation between collimation factor and angular size. To further investigate the relation between R_{coll} and L_{bol} , an analysis was made of all sources whose angle size is five times larger than the beam size. Figure 4 plots the R_{coll} vs. the L_{bol} for these sources. The solid line is the linear least-square fit: $\log(R_{\text{coll}}) = (0.46 \pm 0.04) + (-0.03 \pm 0.01) \log(L_{\text{bol}})$; the correlation coefficient is also poor, $r = -0.28$. The average values of R_{coll} for the sources with large angle sizes are 2.16 with a standard deviation 0.91 for high mass sources and 3.00 with a standard deviation 2.02 for low mass sources. Both are better than those before the effect of beam size was removed (2.05 ± 0.96 and 2.81 ± 2.16 for

the two group sources). It shows again that the collimation degree tends to be lower with increasing bolometric luminosity of the centre source. However, the weak correlation between R_{coll} and L_{bol} remains for sources with large angle sizes, as Fig. 2 shows. This may be related to the driving processes of the outflows. Outflows driven by wide-angle wind may be less collimated than those driven by jets. Now it is believed that low mass outflows are driven by bow shocks (Lee et al. 2000; Chernin & Masson 1995; Raga & Cabrit 1993; de Young 1986) while wide-angle wind may drive massive outflows (Molinari et al. 2002). Low angular resolution also made it difficult to see the fine structure of multi-flows. For example, IRAS 05358+3543, detected as one outflow, was separated into at least three outflows by the Plateau de Bure interferometer (Beuther et al. 2002a,c), including one highly collimated outflow. However, this is not to say that instruments determine the values of collimation. Of the sources depicted in Fig. 3, some small angle size outflows still have good collimation.

The correlation between R_{coll} and L_{bol} is not as obvious as that between outflow mass and bolometric luminosity (see next section). Both Fig. 2 and 4 show that for collimation there is no definite dependance on the bolometric luminosity of the driving sources. Besides the sensitivity and resolutions of the observation equipment, several effects may affect the measurement of collimation degree of the outflows. For example, changing the projection along the line of sight can change the value of the apparent collimation factor for the same bipolar outflow to be smaller or larger than the intrinsic degree of collimation (Wu & Huang 1998). Several factors may influence the intrinsic degree of collimation. Physical conditions of the environment will affect flow morphology. Recently a numerical simulation showed that environmental structure may determine the collimation degree of observed outflows of massive YSOs (Bonnell & Bate 2002). Evidence was also found for the existence of an aligned magnetic field in the outflow regions (Greaves et al. 2001; Houde et al. 2001). Lastly, collimation may also be related to the evolution stage of the central object, and to the acceleration or deceleration phase of the flow itself (Cabrit & Bertout 1986, 1990; Shu et al. 1987). The outflows in Class 0 objects with a cold black body spectral energy distribution (SED) are more collimated than those in Class I objects whose SED is wider than that of a black body and whose positive indices show a more evolved status (Bachiller 1996; Bontemps et al. 1996; Gueth & Guilloteau 1999).

4. Physical parameters

Although the physical parameters of outflows have wide ranges, we believe that statistics derived from a large sample should still reveal general characters and correlations

of the parameters of the outflows and indicate their driving mechanisms. R_{max} ranges from 0.01 to 3.98 pc; mass from 10^{-3} to $10^3 M_{\odot}$; momentum from 10^{-3} to $10^4 M_{\odot} \text{ km/s}$; energy: 10^{38} to 10^{48} ergs; dynamic times t from 4×10^2 to 6×10^5 yr; mechanical luminosity L_m from 10^{-5} to $10^3 L_{\odot}$; force F from 10^{-7} to $10^2 M_{\odot} \text{ km/s/yr}$; mass loss rate $\dot{M}$ from 10^{-9} to $10^{-3} M_{\odot}/\text{yr}$. (Notice that the limits of different parameters may not come from the same source.) The property value ranges are wider than those of the sources measured in a single survey (Bally & Lada 1983; Levreault 1985; Fukui 1989; Ridge & Moore 2001; Zhang et al. 2001; Beuther et al. 2002c); these results from the fact that multiple surveys for different mass objects are included in our sample. Our sample also includes the most massive and the smallest outflows (see Sect. 4.2). However sources with parameters near the limit are rare.

We estimate the maximum possible errors that different considerations in deriving physical parameters may bring. The outflow emission at line wings is often accepted as being optically thin. Nevertheless, some authors deal with it using a derived or assumed optical depth, which may bring a factor of about 5 to the final value of outflow mass, momentum, kinetic energy and so on (Snell et al. 1984; Goldsmith et al. 1984; Shepherd & Churchwell 1996a; Garden et al. 1991; Yu et al. 1999). Different authors used different $[\text{CO}]/[\text{H}_2]$ abundance ratios. Usually this will affect the derived masses by a factor of less than 2 (Ridge & Moore 2001). Occasionally it could be larger, from 2.5×10^{-5} (Rodríguez et al. 1982) to 10^{-4} (Garden et al. 1991). We accept a factor of 4 as the abundance error. Some authors use 1.36 as the mean atomic weight of the mixture of hydrogen and helium (see Garden et al. (1991)), while others only consider pure hydrogen molecular gas (see Snell et al. (1984)). Therefore an uncertainty of 0.36 is introduced. Projection effects affect momentum P and energy E_k by factors of 2 and 3 respectively (Goldsmith et al. 1984). Our experience shows that different velocity ranges (ΔV) of the determined outflow emission add 20% uncertainty to the results. The excitation temperature, T_{ex} , is determined by the relative line intensities of at least two different transitions (Garden et al. 1991). It is usually assumed or defined as the brightness temperature of the CO line peak. T_{ex} ranging from 10 to 15 K for low mass sources (Goldsmith et al. 1984; Snell et al. 1980), and from 30 to 50 K for high mass sources (Shepherd & Churchwell 1996a,b; Beuther et al. 2002c; Wu et al. 2004), causes a maximum uncertainty of about 60%. To investigate how different telescope beam sizes affect the outflow physical parameters, we pick out the sources whose angular diameters are larger than 5 beam sizes of the telescope used. 113 of these sources have available L_{bol} and $\dot{M}$. the linear fit of $\log L_{\text{bol}}$ and $\log \dot{M}$ is: $\log \dot{M} = (0.92 \pm 0.14) + (0.51 \pm 0.04) \log L_{\text{bol}}$, $r = 0.73$.

Comparing the $L_{\text{bol}} - M$ relation here to that of all the 235 sources with available L_{bol} and M , which we present in Sect. 4.1, we find that: The slope of > 5 beam sizes sample (0.51) is a little smaller than that of all (0.56), mainly because the majority of outflows more massive than $10^2 M_{\odot}$ disappeared in the > 5 beam sizes sample due to their greater distances. The correlation coefficients of the > 5 beam sizes sample (0.73) is also a little worse ($r = 0.78$ for all), because the > 5 beam sizes sample is less than half of the whole set. However, the above two deviations are not significant. Thus we are able to conclude that the telescope beam size affects little the outflow physical parameters. We also examine the differences which different CO J transition may cause. The mass derived from CO $J = 1-0$ and $2-1$ are expected to be within a factor of 2 to 3 of the actual values (Margulis & Lada 1985), while this difference mainly comes from the optical depth difference which is already accented in our uncertainties. As a result, the outflow mass, M , has an uncertainty factor of 6.4; the outflow force, F , 6.7; the outflow luminosity, L_{m} , 7.1. The bolometric luminosity, L_{bol} , listed in column 13, primarily derived from far-infrared observations such as IRAS, should be fairly accurate (Lada 1985). Even if there were more than one YSO within the IRAS beam, the emission of the most massive YSO will be the main contribution to the total FIR flux according to the mass-luminosity relation (Casoli et al. 1986). Cabrit & Bertout (1992) listed L_{bol} of 16 outflow sources. We compared these L_{bol} values to those listed in our Table 1a, and found that the uncertainty is within 2~3 except for Orion-A ($(1\sim10)\times10^4$ and 2.1×10^5) and S140 ($(0.5\sim9.0)\times10^4$ and 5×10^3). For most cases, the variation of luminosities provided by different authors in our sample is about 2 to 3, except for a few high mass sources. The errors estimated above are much less than the scatter of the data themselves.

4.1. Relation between the outflow mass and the bolometric luminosity of the center sources

A plot of outflow mass relative to the bolometric luminosity of driving sources shows that the two parameters are physically related. The mass of the outflow is a fundamental parameter. Its relationship to the luminosity of the central source is essential to investigate the nature of the outflows.

Figure 5 plots the mass of outflows M vs. the bolometric luminosities of center sources L_{bol} . It shows that the flow mass increases with the luminosity as a power law. The outflows with higher luminosity values tend to have larger flow masses and vice versa. The least square linear fit in log-log space shown in the figure is $\log M = (-1.04 \pm 0.08) + (0.56 \pm 0.02) \log L_{\text{bol}}$ with the correlation coefficient $r = 0.78$. The strong correlation between outflow mass and luminosity is an interesting result which is the first de-

mostration of such a clear correlation. It presents essential clues to the origin and properties of outflows. The correlation suggests that the outflow mass is linearly correlated with the bolometric luminosity of the center source. The outflow however could not be driven directly by radiation pressure if the photons emitted from the center objects are scattered once before they escape (Bally & Lada 1983, see also Sect. 4.2). This is a classic example in which two parameters are correlated because individually each one is correlated to a third underlying parameter. In our case, it could be that the bolometric luminosity is correlated to the accretion rate (Kenyon et al. 1990; Ohashi et al. 1991; Mundy et al. 1992; Greene et al. 1994), which in turn is correlated to the mass loss rate in the outflow, (Tomisaka 1998; Shu et al. 1994; Contopoulos & Sauty 2001; Fernández & Comerón 2001) and the mass loss rate in the outflow sources is what determines the mass swept up in the outflow lobes (Snell et al. 1980; Ho et al. 1982; Bally & Lada 1983; Goldsmith et al. 1984). So although the radiation energy that the source provided is not able to drive such an outflow mass, they are still correlated.

4.2. Relation between the outflow mechanical luminosity, required force and the bolometric luminosity of the center sources

To investigate the relationship between outflows and their central sources, we further plot the mechanical luminosity (L_{m}) of outflows vs. bolometric luminosity (L_{bol}) of the center stellar objects, shown in Fig. 6. The dashed line shows the equation $L_{\text{m}} = L_{\text{bol}}$. The plot shows that for all sources, L_{m} is less than L_{bol} . The closest point to the dashed line is outflow No. 21, the source L1448 U-star. Bachiller & Cernicharo (1990) pointed out that it was near the upper edge of the populated zone in the mechanical power versus stellar luminosity diagram by Lada (1985). Now that the size of the sample is much larger, the closest point is still located at the upper edge. The lowest point represents outflow No. 337, the source GN21.38.9. This is the lowest-mass Bok globule according to Duvert et al. (1990).

In Fig. 6, the solid line is a linear least-square fit: $\log L_{\text{m}} = (-1.98 \pm 0.14) + (0.62 \pm 0.04) \log L_{\text{bol}}$. The correlation coefficient is 0.69. One can see that the two lines are not parallel. The average deviation between the two lines is larger for the high mass group than the low mass ones. The average values of the ration $L_{\text{m}}/L_{\text{bol}}$ are 0.033 ± 0.086 and 0.0038 ± 0.017 for the low mass and high mass group sources, respectively.

Figure 7 is the plot of the force required to drive the outflow against the bolometric luminosity of the central source. The dashed line shows $F = L_{\text{bol}}/c$. All the outflows are above the line, which means that the radiation pressure of the central sources would not be enough to

drive the outflows if the radiation photons from the central source were scattered once. The two plotted parameters are still correlated. The solid line presents the least square fit for the force of the flow as a function of the bolometric luminosity: $\log F = (-0.92 \pm 0.15) + (0.648 \pm 0.043) \log L_{\text{bol}}$. The correlation coefficient is 0.72.

Bally & Lada (1983) mapped a sample of sources using a single telescope; they discovered a weak correlation between the mechanical luminosity, the required driving force of the outflow and the bolometric luminosity of the central source. The correlation means that the energy and momentum of an outflow is determined by the central luminosity or mass and the physical driving engines are similar for all the sources (Bally & Lada 1983). The above authors also pointed out that the scatter of the plots could be understood as the result of several selection effects. The uncertainties in determining luminosities of central sources can cause significant scattering. A large part of the scatter may be due to the geometrical velocity determination.

To test the correlation between the mechanical luminosity, the force and the luminosity of the central source, the increased sample size is expected to ensure that the scatter in $L_{\text{m}}-L_{\text{bol}}$ plane will be more than that in the $F-L_{\text{bol}}$ plane (Bally & Lada 1983). Here we have a sample 25 times the size of theirs; the correlation coefficient of the line fit for the $F-L_{\text{bol}}$ (0.72) is just slightly better than that for $L_{\text{m}}-L_{\text{bol}}$ (0.69). The standard deviation from the fitted line of $F-L_{\text{bol}}$ is about the same as that of $L_{\text{m}}-L_{\text{bol}}$ or without a significant difference. This may be caused by various uncertainties in our sample, which are mentioned in Sect. 4, while the sample of Bally & Lada (1983) came from a single survey. The parameter ranges of our sample are wider than the sample in Bally & Lada (1983). For example, outflow mass is $10^{-3} \sim 10^3 M_{\odot}$, while theirs is from $0.1 \sim 100 M_{\odot}$.

The mass entrainment rate (M/t) is also correlated linearly to L_{bol} (Wu et al. 2004). This is further evidence showing the dependence of outflow on its driving source.

4.3. Dynamic time

4.3.1. The average age

The dynamic time scale, t , is available for 275 sources. The time scales range from 400 yr (see NGC2024 FIR6, Richer 1990) to 6×10^5 yr (see L1551NE, Moriarty-Schieven & Wannier 1991). The average value is 9.8×10^4 yr and 5.0×10^4 yr for the high mass and low mass group respectively. To investigate why the dynamical timescale of high mass sources appears longer than that of low mass sources, we examined the sources with $t > 2 \times 10^5$ yr. There are 21 such sources, among which 14 belong to the high mass group and 7 belong to the low

mass group. We list all the sources with $t > 2 \times 10^5$ yr in Table 2.

Among 14 high mass sources, 10 are UC HII or bright far infrared sources, which are more evolved among massive YSOs. Another reason why the high mass group has a higher average age is the observation resolution: 14 of the above 21 sources have large distances $D > 1.0$ kpc, while medium (10~15m) or small telescopes (3~5m) were employed to observe 13 of them; among these 13 sources, 12 are high mass sources.

4.3.2. The relation between outflow mass and dynamic time

The plot of the outflow mass against the dynamic time is shown in Fig. 8.

For 98.5% (271/275) of the sources the dynamic timescales are between $10^3 \sim 5.5 \times 10^5$ years, including 99% (155/157) of the low mass sources, and 98% (116/118) of the high mass ones. For the low mass group, the flow mass seems to increase with time while there is no such simple trend for the high mass group. Linear fitting shows $\log M = (1.25 \pm 0.11) + (0.30 \pm 0.12) \log t$, and a poor $r = 0.20$. The correlation is much stronger for the low mass group, $\log M = (-0.71 \pm 0.08) + (0.64 \pm 0.11) \log t$, and $r = 0.41$. Thus the flow processes may be different for the two groups.

5. Associated phenomena

Numerous associated objects were detected and investigated. Ninety (23%) sources have associated HII or centimeter continuum emission regions, 136 (35%) have optical jets or HH objects, 92 (24%) have H_2 jet and 190 (49%) have H_2O masers. Among 136 sources with HH objects or optical characters, 14 (10%) are in the high mass group. In the 190 sources with water masers, 75 (39%) are in the low mass group. There are 45 (12%) outflows with collapses (the percentage in the high mass group is 20%), nearly all of the collapse candidates detected so far. According to Shu et al. (1987), the collapse should occur earlier than bipolar outflow in star formation stages. However, nearly all of the collapses were detected in outflow sources. This may be because the infall material near the embedded sources is shielded by surrounding gas and dust, thus is difficult to observe until the outflow later expels the surrounding material so that the internal region is observable.

6. Spatial distribution and occurrence rate

The three dimensional spatial distribution of the sources projected on the coordinate plane is given in Fig. 9. The sun is at the origin, with the X axis pointing to the 90 degree longitude direction of the IAU Galactic coordinate,

the Y axis passing through the Galactic Center and the Z axis pointing to the north Galactic pole. High mass sources are represented by clear squares and low mass ones by filled triangles. Figure 9a shows the sources projected on the Galactic plane; Fig. 9b is the same frame as Fig. 9a but shows only the inner 2 kilo-parsecs from the sun. Figure 9c and Fig. 9d are projections onto the X-Z and Y-Z planes, respectively.

A prominent feature seen in the plots is that there are more sources between $l = 30^\circ$ to $l = 210^\circ$ than in the rest ($\sim 2 : 1$; see Table 1a). This is largely attributed to the fact that most observations were performed on sources that can be seen in the northern hemisphere. There do exist spatial structures that are asymmetrically distributed with respect to the sun in the local distance, e.g. the neutral interstellar gas (Frisch & York 1983). Such asymmetry mirrors the distribution of the nearby O and B stars associated with the Gould Belt (Stothers & Frogel 1974). The contour of $N(\text{H}) \sim 5 \times 10^{20} \text{cm}^{-2}$ is shown in Fig. 9a as a thick line (Frisch & York 1983). A molecular disk model with radius 1.5kpc and a molecular ring with radius 3-7 kpc are also given in Fig. 9a (Liszt & Burton 1978; Scoville & Sanders 1987). Multiple components along the line of the sight towards these molecular rings where $|l|$ is less than 40° may be another reason that makes it difficult to detect or to identify molecular outflows in these regions.

There are 230 sources in the catalogue within one kilo parsec of the sun. The occurrence rate can be estimated as $74/t \text{ kpc}^{-2} \text{yr}^{-1}$, where t is the average lifetime (in years) of the outflows. For $t = 5.8 \times 10^4$ years, which is the average value of the 230 sources within one kiloparsec, the birth rate is $1.3 \times 10^{-3} \text{ kpc}^{-2} \text{yr}^{-1}$.

Among the 230 sources within 1 kpc of the sun there are 171 sources with available bolometric luminosity (see Table 3). Taking the upper and lower limit of masses of the evolution tracks on the observed H-R diagram according to bolometric luminosity (Levreault 1985, and references therein), the average mass of these sources was estimated to be $3.6M_\odot$ as an upper limit, and $2.1M_\odot$ as a lower limit (see Table 3).

The birth rate measured for the number of stars, $1.3 \times 10^{-3} \text{ kpc}^{-2} \text{yr}^{-1}$, is equal to $4.7 \times 10^{-9} M_\odot \text{pc}^{-2} \text{yr}^{-1}$ measured as a stellar mass upper limit; for the lower limit, it is $2.7 \times 10^{-9} M_\odot \text{pc}^{-2} \text{yr}^{-1}$ (see Table 3). Both are similar to the local star formation rate $3 \sim 6 \times 10^{-9} M_\odot \text{pc}^{-2} \text{yr}^{-1}$, or $4 \sim 11 \times 10^{-9} M_\odot \text{pc}^{-2} \text{yr}^{-1}$, depending on whether there is any compression in the spiral shock wave or not (Kaufman 1979). These results suggest that high-velocity outflows are common in star formation.

7. Summary

In this paper, the searches of molecular outflow sources are reviewed. Our Catalogue includes 391 high velocity molecular outflows and their basic parameters. We classified them into two groups according to the available bolometric luminosity or to the outflow mass. We analyzed the progress of molecular outflow searches and divided the history of outflow search into four periods: I. 1976-1984, II. 1985-1989, III. 1990-1995 and IV. 1996-2003. The characters of different periods and the developmental trend are discussed. The morphology, collimation and physical parameters of outflows were investigated. We also studied the correlation between outflows and their central sources. Finally the spatial distribution and birth rate of young stars are presented and discussed. For convenience, the main statistical results are listed in Table. 4.

Our main results are as follows:

1. The catalogued outflows have been classified into high mass and low mass groups according to the bolometric luminosity of the center sources and the outflow masses. The pace of outflow identification has accelerated in the past two decades. The number of outflows detected per year tends to increase, that is, 7 in Period I but 23 in Period IV. The number of low mass sources increased fast during period II. In period IV, the number of high mass outflows increased rapidly, suggesting that the outflow phenomenon is also common in high mass star formation regions. Data acquired recently suggest high mass star formation processes may also go through mass accreting and ejecting, similar to those processes in low mass star formation.
2. Polarity was known for all the 391 sources, 84% of which are bipolar. Red mono-polar outflows are found to be equally abundant compared to the blue ones within the statistical uncertainty. This suggests that monopolarity of the outflows is intrinsic and is a result of the driving source being partially submerged in the molecular cloud. The number of multiple sources has increased and isolated ones decreased since the last catalogue (Wu et al. 1996), which shows the improvement of the detection technology.
3. We analyzed the collimation of the outflows. The number of sources with known collimation factors has increased from 26 in period I to 213 in period IV. The current average value of the collimation factors is slightly larger than that in period I. Although the resolution of the equipment used in period IV has generally improved, the number of high mass outflows observed increased in this period. As high mass outflows have in general lower collimation factor values, the average collimation value only changed slightly.
4. The collimation of low mass sources is better than that of high mass ones as a whole. Different outflow driving

mechanisms may explain such differences. Moreover, projection effects, physical conditions of the environment and evolutionary states also affect the morphology of outflows.

5. The relationship between physical parameters of the outflow and its central source is examined. Analysis shows that outflow masses are correlated strongly with bolometric luminosities of young stellar sources. This relation, a typical example of an indirect correlation of two quantities, is obtained for the first time.
6. Similar linear correlations exist between flow mechanical luminosity, the force required to drive the outflow and the bolometric luminosity in logarithmic scales. The weak correlation found by Bally & Lada (1983) in an early survey is confirmed.
The mass, momentum and energy of the outflows depend on the luminosity of the central source. These relationships between the parameters of the outflows and the driving sources suggest that the energetic outflows originate from their central source. Although the radiation pressure could not drive the outflow directly if the photons emitted from the central source scatter once, these correlations confirm the relationship between outflows and their central stellar objects.
7. Outflow mass and dynamic time tends to be correlated for low mass sources while for high mass ones there was no such correlation. This together with the differences between the low mass and high mass group sources shown by L_m v.s. L_{bol} (Fig. 6) and F vs. L_{bol} (Fig. 7) is consistent with the presence of different driving mechanisms.
8. Objects associated with outflows were investigated. 136 outflows have associated HH objects; 10% of those are among the high mass group; 190 outflows have water masers, 39% belong to the low mass group. 12% of the outflow sources have ongoing collapse of infall detected.
9. The spatial distribution of known outflows is not symmetric with respect to the sun because the southern hemisphere has fewer identified sources, largely due to less observation. The occurrence rate within 1kpc from the sun is $74/t \text{ kpc}^{-2}\text{yr}^{-1}$, where t is a typical lifetime of outflow in years. For an outflow dynamic time of 5.8×10^4 years the birth rate is $1.3 \times 10^{-3} \text{ kpc}^{-2}\text{yr}^{-1}$, close to the local star birth rate.

Higher angular resolution and sensitivity are necessary to further resolve the velocity structure and survey the distant sources. If a large number of outflows with more accurate morphology are found, we may better understand the physical nature of the energetic processes occurring during early stellar evolution.

Acknowledgements. We would like to thank the anonymous referee for constructive suggestions and helpful comments.

We are very grateful to J. Wu, J. Wang, C. Yang, M. Wei, W. Zhou and L. Zhu for their assistance in the data collection and data checking. We thank Hongping Du and the editor for help with our English expressions. This project is supported by the grants from NSFC No. 10133020, 10128306 and 10203003, and by G1999075405 of NKBRSF.

References

- Bachiller, R. 1996, *ARA&A*, 34, 111
 Bachiller, R. & Cernicharo, J. 1990, *A&A*, 239, 276
 Bachiller, R., Martín-Pintado, J., & Planesas, P. 1991, *A&A*, 251, 639
 Bally, J., Johnstone, D., Joncas, G., Reipurth, B., & Mallén-Ornelas, G. 2001, *AJ*, 122, 1508
 Bally, J. & Lada, C. J. 1983, *ApJ*, 265, 824
 Beuther, H., Schilke, P., Gueth, F., et al. 2002a, *A&A*, 387, 931
 Beuther, H., Schilke, P., Menten, K. M., et al. 2002b, *ApJ*, 566, 945
 Beuther, H., Schilke, P., Sridharan, T. K., et al. 2002c, *A&A*, 383, 892
 Bonnell, I. A. & Bate, M. R. 2002, *MNRAS*, 336, 659
 Bontemps, S., André, P., Terebey, S., & Cabrit, S. 1996, *A&A*, 331, 858
 Cabrit, S. & Bertout, C. 1986, *A&A*, 307, 313
 Cabrit, S. & Bertout, C. 1990, *A&A*, 348, 530
 Cabrit, S. & Bertout, C. 1992, *A&A*, 261, 274
 Casoli, F., Combes, F., Dupraz, C., Gerin, M., & Boulanger, F. 1986, *A&A*, 169, 281
 Chernin, L. & Masson, C. R. 1995, *ApJ*, 455, 182
 Choi, M., Evans, N. J. I., & Jaffe, D. T. 1993, *ApJ*, 417, 624
 Contopoulos, I. & Sauty, C. 2001, *A&A*, 365, 165
 de Young, J. S. 1986, *ApJ*, 307, 62
 Duvert, G., J. Cernicharo, Bachiller, R., & Gómez-González, J. 1990, *A&A*, 233, 190
 Evans, N. J. 1999, *ARA&A*, 37, 311
 Felli, M., Palagi, F., & Tofani, G. 1992, *A&A*, 255, 293
 Fernández, M. & Comerón, F. 2001, *A&A*, 380, 264
 Frisch, P. C. & York, D. G. 1983, *ApJ*, 271, 59
 Fuente, A., Neri, R., Martín-Pintado, J., et al. 2001, *A&A*, 366, 873
 Fukui, Y. 1989, *ESO Workshop on Low Mass Star Formation and Pre-Main Sequence Objects*, ed. B. Reipurth, 95
 Garay, G., Kohnenkamp, I., Bourke, T. L., guez, L. F. R., & Lehtinen, K. K. 1998, *ApJ*, 509, 768
 Garden, R. P., Hayashi, M., Hasegawa, T., Gatley, I., & Kaifu, N. 1991, *ApJ*, 374, 540
 Goldsmith, P. F., Snell, R. L., Hemeon-Heyer, M., & Langer, W. D. 1984, *ApJ*, 286, 599
 Greaves, J. S., Holland, W. S., & Ward-Thompson, D. 2001, *ApJ*, 546, L53

- Greene, T. P., Wilking, B. A., André, P., Young, E. T., & Lada, C. J. 1994, *ApJ*, 434, 614
- Gueth, F. & Guilloteau, S. 1999, *A&A*, 343, 571
- Herbig, G. 1960, *ApJS*, 337, 4
- Ho, P. T. P., Moran, J. M., & Rodríguez, L. F. 1982, *ApJ*, 262, 619
- Hofner, P., Wiesemeyer, H., & Henning, T. 2001, *ApJ*, 549, 425
- Houde, M., Thomas, P. G., Pierre, B., Ruisheng, P., & Hiroshige, Y. 2001, *ApJ*, 547, 311
- Kaufman, M. 1979, *ApJ*, 232, 707
- Kenyon, S. J., Hartmann, L. W., Strom, K. M., & Strom, S. E. 1990, *AJ*, 99, 869
- Koo, B. C. 1989, *ApJ*, 337, 318
- Kroupa, P., Tout, C. A., & Gilmore, G. 1990, *MNRAS*, 244, 76
- Kuhi, L. V. 1964, *ApJ*, 140, 1409
- Kwan, J. & Scoville, N. 1976, *ApJ*, 210, L39
- Lada, C. J. 1985, *ARA&A*, 23, 267
- Lane, A. P. 1989, *ESO Workshop on Low Mass Star Formation and Pre-Main Sequence Objects*, ed. B. Reipurth, 89
- Lee, C., Mundy, L. G., Reipurth, B., Ostriker, E. C., & Stone, J. M. 2000, *ApJ*, 542, 925
- Levreault, R. M. 1985, *Molecular Outflows and Mass Loss in Pre-Main Sequence Stars*, Tech. Rep. 85-1, Astronomy Dept. Univ. of Texas
- Levreault, R. M. 1988, *ApJS*, 67, 283
- Liszt, H. S. & Burton, W. B. 1978, *ApJ*, 226, 790
- Lizano, S., Heiles, C., Rodríguez, L. F., et al. 1988, *ApJ*, 328, 763
- Margulis, M. & Lada, C. J. 1985, *ApJ*, 299, 925
- Matzner, C. D. & Mckee, C. F. 1999, *ApJ*, 526, 109
- Megeath, S. T. & Tiefertunk, A. R. 1999, *ApJ*, 526, L113
- Molinari, S., Testi, L., Rodríguez, L. F., & Zhang, Q. 2002, *ApJ*, 570, 758
- Moriarty-Schieven, G. H. & Wannier, P. G. 1991, *ApJ*, 373, L23
- Mundy, L. G., Wootten, A., Wilking, B. A., Blake, G. A., & Sargent, A. I. 1992, *ApJ*, 385, 306
- Myers, P. C., Mardones, D., Tafalla, M., Williams, J. P., & Wilner, D. J. 1996, *ApJ*, 465, 133
- Narayanan, G. & Walker, C. K. 1996, *ApJ*, 466, 884
- Natta, A., Giovanardi, C., Palla, F., & Evans, N. J. 1988, *ApJ*, 327, 817
- Ohashi, N., Kawabe, R., Hayashi, M., & Ishiguro, M. 1991, *AJ*, 102, 2054
- Raga, A. C. & Cabrit, S. 1993, *A&A*, 278, 267
- Reipurth, B., Heathcote, S., Morse, J., Hartigan, P., & Bally, J. 2002, *AJ*, 123, 362
- Richer, J. S. 1990, *MNRAS*, 245, 24P
- Ridge, N. A. & Moore, T. J. T. 2001, *A&A*, 378, 495
- Rodríguez, L. F., Carral, P., Moran, J. M., & Ho, P. T. P. 1982, *ApJ*, 260, 635
- Russell, A. P. G., Bally, J., Padman, R., & Hills, R. E. 1992, *ApJ*, 387, 219
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Scoville, N. Z. & Sanders, D. B. 1987, *H₂ in the Galaxy, in Interstellar Processes*, ed. D. J. Hollenbach & H. A. Thronson (Jr, D. Reidel Publishing Com.), 21
- Shepherd, D. S. & Churchwell, E. 1996a, *ApJ*, 472, 225
- Shepherd, D. S. & Churchwell, E. 1996b, *ApJ*, 457, 267
- Shepherd, D. S., Watson, A. M., Sargent, A. I., & Churchwell, E. 1998, *ApJ*, 507, 861
- Shu, F., Najita, J., Ostriker, E., et al. 1994, *ApJ*, 429, 781
- Shu, F. H., Adams, F. C., & Lizano, S. 1987, *ARA&A*, 25, 23
- Snell, R. L., Loren, R. B., & Plambeck, R. L. 1980, *ApJ*, 239, L17
- Snell, R. L., Scoville, N. Z., Sanders, D. B., & Erickson, N. R. 1984, *ApJ*, 284, 176
- Stothers, R. & Frogel, J. A. 1974, *AJ*, 79, 456
- Tomisaka, K. 1998, *ApJ*, 502, L163
- Wolf-Chase, G. A., Barsony, M., Wootten, H. A., et al. 1998, *ApJ*, 501, L193
- Wu, Y. & Huang, M. 1998, *Chin. Phys. Lett*, 15, 388
- Wu, Y., Huang, M., & He, J. 1996, *A&AS*, 115, 283
- Wu, Y., Wu, J., & Wang, J. 2001, *A&A*, 380, 665
- Wu, Y., Yang, C., Li, Y., et al. 1999, *Science in China (Ser. A)*, 42, 732
- Wu, Y., Zhang, Q., Chen, H., et al. 2004, submitted by *AJ*
- Yu, K., Billawala, Y., & Bally, J. 1999, *AJ*, 118, 2940
- Zhang, Q., Hunter, T. R., Brand, J., et al. 2001, *ApJ*, 552, L167
- Zhou, S. 1992, *ApJ*, 394, 204
- Zuckerman, B., Kuiper, T. B. H., & Kuiper, E. N. R. 1976, *ApJ*, 209, 137
- Zuckerman, B. & Palmer, P. 1975, *ApJ*, 199, 35

Table 2. Outflow with dynamic time $> 20 \times 10^4$ yr

Outflow No.	Outflow Name	D kpc	t 10^4 yr	Group	Telescope used	Ref.
009	00494+5617	2.2	24	H	14m	3
014	02310+6133	3.8	51.5	H	13.7m	381
015	02461+6147	4.2	35	H	3m($J = 3 - 2$)	380
040	B5-IRS4	0.35	24	L	7m	21
052	L1551NE	0.16	60	L	12m	128
126	05490+2658	2.1	37	H	14m	3
130	05553+1631	2.5	46	H	14m	3
	G192.16	2.0	27	H	12m	194
131	CB39	0.6	30	L	14m	178
139	S255	2.5	50	H	14m	117
162	07028-1100	1.4	24	L	13.7m	421
166	WB89 1135	6.42	37.9	H	15m	196
173	WB89 1275	6.38	27.8	H	15m	196
237	18150-2016	1.8	24	H	4m	260
281	19471+2641	2.62	21	H	10m($J = 3 - 2$)	203
325	21015+6757	0.44	24	L	3m($J = 3 - 2$)	397
336	V645 Cyg	5.6	50	H	5m($J = 2 - 1$)	8
337	GN21.38.9	0.75	30	L	10m	150
350	21432+4719	1.0	22	L	45m	258
360	22142+5206	4.5	23	H	45m	187
374	22475+5939	4.7	29	H	12m	199
382	MBM55	0.18	22	H	12m	100

Table 3. L_{bol} and mass ranges for associated stellar sources of outflow within 1 kpc.

$L_{\text{bol}}(L_{\odot})$	$< 10^{-1}$	$10^{-1} \sim 10^0$	$10^0 \sim 10^1$	$10^1 \sim 10^2$	$10^2 \sim 10^3$	$10^3 \sim 10^4$	$> 10^4$
Source Number	0	20	50	49	36	8	8
Upper limit $M_{\odot}$	1.0	1	1.5	3	5	9	15
lower limit $M_{\odot}$	0.2	0.5	1.0	1.5	3	5	9

Table 4. Index of main statistical results

Item	Result	Section	Paragraph
Outflows cataloged	391+6	2.1	2
High group ($L_{\text{bol}} > 10^3 L_{\odot}$ or $M > 3M_{\odot}$)	139	2.3	2
low group ($L_{\text{bol}} \leq 10^3 L_{\odot}$ or $M \leq 3M_{\odot}$)	223	2.3	2
Sources detected in Period I, II, III, IV	55, 95, 88, 159	2.4	2
High group in Period I, IV	31%, 38%	2.4	3
Sources with $L_{\text{bol}} > 10^4$ to $L_{\odot}$	23%	2.4	4
Sources with UC HII regions	23%	2.4	4
Bipolar sources	327(84%)	3	1
Monopolar sources	50(13%) [28 red, 22 blue]	3	1
Multipolar sources	12(3%)	3	1
Isotropic sources	2	3	1
Sources with available R_{coll}	213	3	3
Average R_{coll}	2.45 ± 1.74	3	3
Average R_{coll} of high group	2.05 ± 0.96	3	3
Average R_{coll} of low group	2.81 ± 2.16	3	3
Bolometric luminosity range	$0.1\text{-}10^6 L_{\odot}$	2.3	1
Uncertain factor of M	6.4	4	2
Uncertain factor of F	6.7	4	2
Uncertain factor of L_{m}	7.1	4	2
R_{max} range	0.01 to 3.98 pc	4	1
Mass range	10^{-3} to $10^3 M_{\odot}$	4	1
Momentum	10^{-3} to $10^4 M_{\odot} \text{km/s}$	4	1
Energy	10^{38} to 10^{48} ergs	4	1
Mechanical luminosity	10^{-5} to $10^3 L_{\odot}$	4	1
Force F	$10^{-7} \sim 10^2 M_{\odot} \text{km/s yr}$	4	1
Mass loss rate $\dot{M}$	$10^{-9} \sim 10^{-3} M_{\odot}/\text{yr}$	4	1
Dynamical time available sources	275	4.3.1	1
Range of dynamical time	4×10^2 to 6×10^5 yr	4.3.1	1
Dynamical time of the 99% sources	$10^3 \sim 5.5 \times 10^5$ yr	4.3.2	2
Average dynamical time of high group	9.8×10^4 yr	4.3.1	1
Average dynamical time of low group	5.0×10^4 yr	4.3.1	1
Sources with $t > 2 \times 10^5$ yr	21 [14 in high group]	4.3.1	1
Sources associated with HH objects	136 (35%) [90% in low group]	5	
Sources associated with H_2 jet	92 (24%)	5	
Sources associated with H_2O masers	190 (49%) [115 (61%) in high group]	5	
Collapse candidates	45 (12%)	5	
Sources within 1 kpc	230	6	3
Sources within 1 kpc (L_{bol} available)	171 [16(9%) in high group]	6	4




Fig. 1. Frequency distributions of detected outflow numbers with time. a). Histogram of the mapped outflow numbers in the four periods; the widths of the shadow zones are determined by how long the corresponding period lasting; b). Histogram of average numbers per year in the four periods.




Fig. 2. Collimation factors versus bolometric luminosity of the associated infrared source (L_{bol}). The filled triangles represent the sources from the low mass group, while the open boxes indicate the sources from the high mass group. The solid line is the least square linear fit line.




Fig. 3. Collimation factors versus angular sizes. The symbols are the same as for Fig. 2.




Fig. 4. Collimation factors versus bolometric luminosity (L_{bol}) for sources with angular sizes at least five times the beam size. The symbols are the same as for Fig. 2.




Fig. 5. The outflow mass versus the bolometric luminosity (L_{bol}). The symbols are the same as in Fig. 2. The solid line is the least square linear fit.




Fig. 6. The outflow luminosity (L_m) versus bolometric luminosity (L_{bol}). The symbols are the same as for Fig. 2. The relation $L_m = L_{\text{bol}}$ is shown as a dashed line. The solid line is the least square linear fit line.




Fig. 7. Outflow force F versus bolometric luminosity (L_{bol}) of the associated infrared sources. The symbols are the same as in Fig. 2. The dashed line presents the relation $F = L_{\text{bol}}/c$. The solid line is the least square linear fit.




Fig. 8. Outflow masses M versus dynamical time. The symbols are the same as for Fig. 2. The dashed and solid lines are the least square linear fit for sources of the high and low mass group respectively.





Fig. 9. Spatial distribution of outflows. The sun is at the origin of the coordinate. The X-axis points to the 90° longitude direction of the IAU Galactic coordinate, the Y-axis passes through Galactic Center and the Z-axis points to the north Galactic pole. Triangles represent low mass outflows and circles high mass ones. The thick dark grey line within the 2 kpc radius circle denotes the contour of $N(\text{H}) \sim 5 \times 10^{20} \text{ cm}^{-2}$ (Frisch & York 1983). a). The outflow distribution on the coordinate plane x-y. The three circles from inner to outer indicate distances of 2 kpc, 5 kpc and 10 kpc from the sun. The small shadow elliptic presents the projection of a molecular disk model with a radius of 1.5 kpc (Liszt & Burton 1978). The shadow ring indicates projection of the molecular ring with radius 3~7 kpc (Scoville & Sanders 1987). b). The same as a), but shows only the inner 2 kpc from the sun. The two circles indicate 1 kpc and 2 kpc from the sun. c). Projection onto the X-Z plane. d). Projection onto the Y-Z plane. Note that the vertical axes in c) and d) are scaled up for clarity.

**Fig. 9.** *continued*